

# Difficult Airway due to Retropharyngeal Hematoma after Stabbing to the Neck

Kouhei Iwashita, MD<sup>1\*</sup>, Kenji Shigematsu, MD<sup>1</sup>, Shinjiro Shono, MD<sup>2</sup>, Keiichi Nitahara, MD, PhD<sup>1</sup> and Kazuo Higa, MD<sup>1</sup>

<sup>1</sup>Department of Anesthesiology, Fukuoka University Faculty of Medicine, Fukuoka, Japan

<sup>2</sup>Department anesthesiology, Chikushi hospital, Fukuoka University, Chikushino, Japan

## Abstract

Reports of retropharyngeal hematoma have been scarce in the anesthesiology literature. We report a patient whose trachea was difficult to intubate due to retropharyngeal hematoma after stabbing to the neck. A woman with a knife injury to the common carotid artery required emergency carotid arterioplasty. When tracheal intubation was attempted, marked swelling of the posterior pharyngeal wall made the vocal cords impossible to visualize. Preoperative computed tomography showed a retropharyngeal hematoma. The patient required mechanical ventilation for 2 days.

**Keywords** Difficult airway; Retropharyngeal hematoma; Stabbing; Carotid artery

**\*Corresponding author:** Kouhei Iwashita, MD, Assistant Professor, Department of Anesthesiology, Fukuoka University Faculty of Medicine, 7-45-1 Nanakuma, Jonan-ku, Fukuoka 814-0180, Japan, E-mail: iwashita@fukuoka-u.ac.jp

**Received Date:** 31 March 2014

**Accepted Date:** 18 July 2014

**Published Date:** 22 July 2014

**Citation:** Iwashita K, Shigematsu K, Shono S, Nitahara K, Higa K (2014) Difficult Airway due to Retropharyngeal Hematoma after Stabbing to the Neck. Enliven: J Anesthesiol Crit Care Med 1(2): 010.

**Copyright:** © 2014 Dr. Kouhei Iwashita. This is an Open Access article published and distributed under the terms of the Creative Commons Attribution License, that permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

## Introduction

Any space occupying lesion in the retropharyngeal space, including a collection of blood, can obstruct the upper airway [1]. Retropharyngeal hematoma (RPH) is rare; however, it can obstruct the airway rapidly [2-5], and can be fatal [5-7]. When dyspnea or stridor occurs in RPH, tracheal intubation is mandatory to secure the airway [8]. However, tracheal intubation is technically difficult in some patients with RPH [5,8,9]. Retropharyngeal hematoma caused by stabbing to the neck has not previously been reported.

We report difficult tracheal intubation in a patient who had RPH due to common carotid artery injury after stabbing to the neck.

## Case report

A 31 year old woman was stabbed with a knife in the left side of the neck, left chest and upper abdomen. The time of injury was unclear; however, the emergency medical services had been called 2 hours and 40 minutes before the patient was brought to the operating room, having taken 10 minutes to arrive at the scene. At the scene, her blood pressure had been 143/84 mmHg, heart rate 148 beats/min, and respiratory rate 30 breaths/min. Manual compression had been applied to the neck and the patient was transported to our hospital.

On arrival in the emergency department, her blood pressure had fallen to 82/32 mmHg, and her heart rate and respiratory rate had risen to 150 beats/min and 36 breaths/min, respectively. Peripheral blood oxygen saturation (SpO<sub>2</sub>) was 100% with oxygen supplied at 10 L/min by face mask. Manual compression of the wound was released as there was no active bleeding from the neck wound.

Intravenous fluid resuscitation was undertaken, using 2,500 mL crystalloid fluid, 1,000 mL 5% albumin, and 10 units of packed red blood cells. The blood pressure rose to 125/65 mmHg, and the heart rate fell to 105 beats/min. Laboratory investigations performed 1 hour after arrival showed a blood hemoglobin concentration of 11.5 g/dL, and a hematocrit of 33.4%. As blood pressure rose, pulsatile bleeding was observed from the left neck, so manual compression was resumed. Auscultation of the chest was unremarkable. Computed tomography (CT) of the neck revealed a hematoma and traumatic emphysema around the left carotid artery, and deviation of the trachea to the right. A lateral X-ray of the neck was not taken. The chest and abdominal CT revealed intra-abdominal hemorrhage. Emergency arterioplasty of the left carotid artery and abdominal surgery for hemostasis was planned.

She was transferred to the operating room with her neck wound under manual compression. Heart rate, electrocardiogram, SpO<sub>2</sub>, and invasive blood pressure were monitored. There was no stridor and she was able to communicate, speaking without hoarseness. At the time of entering the operating room, the blood pressure was 130/70 mmHg, heart rate 110 beats/min, respiratory rate 20 breaths/min and SpO<sub>2</sub> 100% with oxygen provided at 10 L/min by face mask.

Anesthesia was induced with intravenous propofol 50 mg after 100% oxygen had been administered for 5 minutes by face mask. Manual ventilation was straightforward. Neuromuscular block was achieved with intravenous rocuronium 50 mg. When a second-year grade anesthesiologist tried to intubate the trachea with a Macintosh laryngoscope, the tip of the epiglottis could be visualized, but not the vocal cords, which were obscured by a swelling of the posterior wall of the pharynx that was compressing the epiglottis. Blind tracheal intubation with a stylet was attempted, resulting in esophageal intubation. Additional intravenous propofol 50 mg was administered and a Pentax-Airway Scope® videolaryngoscope (Pentax Co., Tokyo, Japan) was prepared. Meanwhile, an attending anesthesiologist tried tracheal intubation using the stylet. He could also identify the tip of the epiglottis, and although the vocal cords could still not be visualized, he was able to blindly intubate the trachea using the stylet. Expiratory carbon dioxide was detected. Fiberoptic bronchoscopy confirmed that the tube was in the tracheal lumen above the carina. It took 5 minutes to intubate the trachea from induction of anesthesia: SpO<sub>2</sub> had remained 100% throughout. Thereafter, anesthesia was maintained with sevoflurane and fentanyl. After the airway was secure, we reviewed the preoperative CT of the neck (Figure 1). The CT showed that the distance from the ventral line of the cervical spine to the posterior wall of the pharynx was 16 mm at C<sub>2</sub>, 18 mm at C<sub>3</sub>, and 19 mm at C<sub>4</sub>, and that the distance from the ventral line of the cervical spine to the posterior wall of the trachea was 21 mm at C<sub>5</sub>, and 19 mm at C<sub>6</sub>. The vocal cords were situated at the level of C<sub>5</sub>.

Left carotid arterioplasty, and repairs of the diaphragm and gastric serosa were performed. Surgery lasted 4 hours 45 minutes. The patient was mechanically ventilated postoperatively. On postoperative day 1, repeat CT of the neck showed tracheal deviation to the right. There was no swelling of the tracheal mucosa. She was extubated 2 days after surgery. Immediately after extubation, the respiratory pattern was normal, the respiratory rate was 20 breaths/min, and SpO<sub>2</sub> was 100% with oxygen 6 L/min by mask. She experienced transient hoarseness, but there was no inspiratory stridor. Supplemental oxygen was no longer required by the fourth postoperative day. SpO<sub>2</sub> ranged between 98 and 99%. She started oral intake on postoperative day 5 and her recovery was rapid and uneventful. Institutional Review Board of the Fukuoka University Hospital permission to publish this report has been obtained (No. 13-3-21).

## Discussion

Retropharyngeal hematoma can cause life-threatening airway obstruction [5-7]. Fatalities due to RPH have been reported [7]. Reports of RPH have been scarce in the anesthesiology literature, although it may be associated with a variety of conditions, such as cervical spine fracture [10,11], trauma to the head or neck [12,13], foreign body ingestion [14], violent cough or sneeze [15,16], carotid sinus massage [3], coagulopathy [17], anticoagulation therapy [4,18], mycotic aneurysm of the external carotid artery [19], thyroid cyst [1], and stellate ganglion block [8].




Figure 1. Contrast-enhanced computed tomography of the neck. Horizontal view. The patient was supine with her head turned to the right. This slice was taken at the upper level of the vocal cords. Red arrow indicates the distance from the ventral line of the cervical spine to the posterior wall of the pharynx. The black arrows indicate the extent of the retropharyngeal hematoma. Arrow heads indicate the hematoma and emphysema around the wound.

Retropharyngeal hematoma develops in the retropharyngeal space, which is bounded by the buccopharyngeal fascia ventrally, the prevertebral fascia dorsally and the carotid sheaths laterally. It extends caudally from the base of the skull to the posterior mediastinum at the level of T<sub>2</sub> to T<sub>6</sub> [20]. The retropharyngeal space is continuous with the parapharyngeal space [1], and normally contains loose connective tissue, lymph nodes, and fat [11,21]. The parapharyngeal space lies laterally to the pharynx and contains muscles, the carotid sheath, the vagus nerve, loose connective tissue, and fat [1]. In our patient, blood hemorrhaging from the carotid artery appears to have entered the retropharyngeal space, causing RPH.

Lateral cervical spine X-rays and CT images can provide valuable diagnostic and clinical information in the presence of RPH [12]. The maximum normal distance from the ventral line of the cervical vertebral bodies to the posterior wall of the trachea on the lateral cervical spine X-ray is 8-11 mm at C<sub>1</sub>, 6 mm at C<sub>2</sub>, 7 mm at C<sub>3</sub>, 8 mm at C<sub>4</sub>, 22 mm at C<sub>5</sub>, 19-20 mm at C<sub>6</sub>, and 20-21 mm at C<sub>7</sub> [11]. Increases in these distances indicate swelling of the retropharyngeal space. Computed tomography of the neck assists with further diagnosis, as it can help identify the nature of the swelling (including the presence of blood), and quantify lateral swelling, tracheal shift [5], and swelling of the tracheal mucosa [22]. Retropharyngeal hematoma distorts the normal anatomy of the larynx, making tracheal intubation difficult. In our patient, the CT of the neck revealed RPH from C<sub>2</sub> to C<sub>4</sub>. When tracheal intubation was attempted, we could not see the vocal cords because of ventral swelling of the posterior pharyngeal wall. We could have predicted a difficult intubation from the CT images.

Symptoms and signs of RPH include sore throat, neck pain, neck swelling, dyspnea, hoarseness, and stridor. Ecchymosis can appear on the neck or anterior chest wall in some cases [8,12]. Airway obstruction in RPH is thought to be caused by swelling of the pharyngolarynx due to venous and lymphatic congestion [23,24]. The airway should be secured immediately if inspiratory stridor occurs in patients with RPH [8]. This was absent in our patient.

Massive swelling of the pharyngolarynx distorts the normal anatomy of the larynx and makes direct visualization of the vocal cords difficult. Tracheal intubation may be extremely difficult or impossible in some patients with RPH, even for experienced anesthesiologists [5,23]. Although the attendant anesthesiologist was able to intubate the trachea blindly using a stylet in our patient, the vocal cords could not be identified due to the massive protrusion of the posterior pharyngeal wall. When tracheal intubation with a laryngoscope is not possible in patients with RPH, alternative airway management techniques should be adopted. These include fiberoptic endotracheal intubation [10,12,13], the Pentax- Airway Scope® [25], percutaneous transtracheal jet ventilation [26], cricothyroidotomy [6,9], and tracheostomy [5,6,13,27]. Percutaneous transtracheal jet ventilation through the cricothyroid membrane can provide immediate oxygenation in a 'can't ventilate, can't intubate' scenario, with the additional benefit that the high intra-tracheal pressure from the jet ventilation may open the collapsed vocal cords and airway.

Here we report the case of a patient with RPH whose trachea was difficult to intubate. It is important to suspect RPH in patients with carotid arterial injury and hemorrhage. A range of alternative airway devices with which to intubate the trachea should be at hand before induction of anesthesia for patients with RPH.

## References

1. Paleri V, Maroju RS, Ali MS, Ruckley RW (2002) Spontaneous retro-and parapharyngeal haematoma caused by intrathyroid bleed. *J Laryngol Otol* 116: 854-858.
2. Coleman JA Jr, Johnson JT (1986) Retropharyngeal hematoma: complication of cervical fracture. *Otolaryngol Head Neck Surg* 94: 387-390.
3. Thomás MD, Torres A, García-Polo J, Gavilán C (1991) Life-threatening cervico-mediastinal haematoma after carotid sinus massage. *J Laryngol Otol* 105: 381-383.
4. Chiti-Batelli S, Vaz F, Coman S (2005) Traumatic retropharyngeal haematoma in an anticoagulated patient: case report and proposal for a clinical protocol. *Acta Otolaryngol* 125: 443-445.
5. Birkholz T, Kröber S, Knorr C, Schiele A, Bumm K, et al. (2010) A retropharyngeal-mediastinal hematoma with supraglottic and tracheal obstruction: The role of multidisciplinary airway management. *J Emerg Trauma Shock* 3: 409-411.
6. Kuhn JE, Graziano GP (1991) Airway compromise as a result of retropharyngeal hematoma following cervical spine injury. *J Spinal Disord* 4: 264-269.
7. Kashiwagi M, Ikeda N, Tsuji A, Kudo K (1991) Sudden unexpected death following stellate ganglion block. *Leg Med (Tokyo)* 1: 262-265.
8. Higa K, Hirata K, Hirota K, Nitahara K, Shono S (2006) Retropharyngeal hematoma after stellate ganglion block: analysis of 27 patients reported in the literature. *Anesthesiology* 105: 1238-1245.
9. McLaughlan CA, Pidsley R, Vandenberg PJ (1991) Minor trauma - major problem. Neck injuries, retropharyngeal haematoma and emergency airway management. *Arch Emerg Med* 8: 135-139.
10. Lazott LW, Ponzo JA, Puana RB, Artz KS, Ciceri DP, et al. (2007) Severe upper airway obstruction due to delayed retropharyngeal hematoma formation following blunt cervical trauma. *BMC Anesthesiol* 7: 2.
11. Penning L (1981) Prevertebral hematoma in cervical spine injury: incidence and etiologic significance. *AJR Am J Roentgenol* 136: 553-561.
12. Lin JY, Wang CH, Huang TW (2007) Traumatic retropharyngeal hematoma: case report. *Auris Nasus Larynx* 34: 423-425.
13. Kochilas X, Ali A, Montague ML, Kelleher RJ (2004) Retropharyngeal space swelling secondary to minor blunt head and neck trauma. *J Laryngol Otol* 118: 465-467.
14. Ophir D, Bartal N (1988) Retropharyngeal hematoma following fish bone ingestion. *Ear Nose Throat J* 67: 528-530.
15. Genovesi MG, Simmons DH (1975) Airway obstruction due to spontaneous retropharyngeal hemorrhage. *Chest* 68: 840-842.
16. Owens DE, Calcaterra TC, Aarstad RA (1975) Retropharyngeal hematoma. A complication of therapy with anticoagulants. *Arch Otolaryngol* 101: 565-568.
17. Harper M, Obolensky L, Roberts P, Mercer M (2007) A case of acute upper and lower airway obstruction due to retropharyngeal haemorrhage secondary to acquired haemophilia A. *Anaesthesia* 62: 627-630.
18. Lepore ML (1976) Upper airway obstruction induced by warfarin sodium. *Arch Otolaryngol* 102: 505-506.
19. Weaver DF, Young DJ (1964) Retropharyngeal haemorrhage from the carotid arteries. *Ann Otol Rhinol Laryngol* 73: 541-544.
20. Wong YK, Novotny GM (1978) Retropharyngeal space - a review of anatomy, pathology, and clinical presentation. *J Otolaryngol* 7: 528-536.
21. Davis WL, Harnsberger HR, Smoker WR, Watanabe AS (1990) Retropharyngeal space: evaluation of normal anatomy and diseases with CT and MR imaging. *Radiology* 174: 59-64.
22. Carmichael FJ, McGuire GP, Wong DT, Crofts S, Sharma S, et al. (1996) Computed tomographic analysis of airway dimensions after carotid endarterectomy. *Anesth Analg* 83: 12-17.
23. Wade JS (1980) Respiratory obstruction in thyroid surgery. *Ann R Coll Surg Engl* 62: 15-24.
24. Wells DG, Zelcer J, Wells GR, Sherman GP (1988) A theoretical mechanism for massive supraglottic swelling following carotid endarterectomy. *Aust N Z J Surg* 58: 979-981.
25. Asai T (2010) Pentax-AWS videolaryngoscope for awake nasal intubation in patients with unstable necks. *Br J Anaesth* 104: 108-111.
26. Patel RG (1999) Percutaneous transtracheal jet ventilation: a safe, quick, and temporary way to provide oxygenation and ventilation when conventional methods are unsuccessful. *Chest* 116: 1689-1694.
27. El Kettani C, Badaoui R, Lesoin FX, Le Gars D, Ossart M (2002) Traumatic retropharyngeal hematoma necessitating emergency intubation. *Anesthesiology* 97: 1645-1646.

Submit your manuscript at  
<http://enlivenarchive.org/submit-manuscript.php>  
 New initiative of Enliven Archive

Apart from providing HTML, PDF versions; we also provide **video version** and deposit the videos in about 15 freely accessible social network sites that promote videos which in turn will aid in rapid circulation of articles published with us.