

Research Article

www.enlivenarchive.org

Enliven: Microbes and Microbial Techniques

ISSN: 2576-2540

Extended ɛ34 Phage TSP Renatures after Urea-Acid Unfolding

Joseph A. Ayariga^{1*}, Logan Gildea², and Robert Villafane³

¹The Biomedical Engineering Program, College of Science, Technology, Engineering and Mathematics (C-STEM), Alabama State University, 1627 Hall Street, Montgomery, AL, 36104.

²The Microbiology Program, College of Science, Technology, Engineering and Mathematics (C-STEM), Alabama State University, Montgomery, AL, 36104 ³Department of Biological Sciences, Microbiology PhD. Program, College of Science, Technology, Engineering and Mathematics (C-STEM), Alabama State University, 1627 Hall Street Montgomery, Alabama 36104

*Corresponding author: Joseph A. Ayariga, The Biomedical Engineering Program, College of Science, Technology, Engineering and Mathematics (C-STEM), Alabama State University, 1627 Hall Street, Montgomery, AL, 36104, Email: jayariga7546@myasu.alasu. edu

Received Date: 02nd December 2021 Accepted Date: 05th January 2022 Published Date: 10th January 2022 **Citation**: Ayariga JA, Gildea L, Villafane R. Extended ɛ34 Phage TSP Renatures after Urea-Acid Unfolding. Enliven: Microb Microbial Tech. 2022; 9(1): 001.

Copyright: Joseph A. Ayariga @ 2022. This is an Open Access article published and distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Abstract

In antimicrobial-peptide/protein engineering, understanding the peptide/protein's adaptability to harsh environmental conditions such as urea, proteases, fluctuating temperatures, high salts provide enormous insight into the pharmacokinetics and pharmacodynamics of the engineered peptide/protein and its ability to survive the harsh internal environment of the human body such as the gut or the harsh external environment to which they are applied. A previous work in our laboratory demonstrated that our cloned $E\varepsilon34$ TSP showed potent antimicrobial activity against *Salmonella newington*, and more so, could prevent biofilm formation on decellularized tissue. In this work, the effects of urea-acid on the $E\varepsilon34$ stability is studied, and the results demonstrates that at lower pHs of 3 and 4 with urea the protein was denatured into monomeric species. However, the protein withstood urea denaturation above pH of 5 and thus remained as trimeric protein. The mechanism of denaturation of $E\varepsilon34$ TSP seems to show that urea denatures proteins by depleting hydrophobic core of the protein by directly binding to the amide units via hydrogen bonds. The results of our *in-silico* investigation determined that urea binds with $E\varepsilon34$ TSP with relative free energies range of -3.4 to -2.9 kcal/mol at the putative globular head binding domain of the protein. The urea molecules interacts with with the protein's predicted hydrophobic core, thus, disrupting and exposing the shielded hydrophobic moieties of $E\varepsilon34$ TSP to the solvent. We further showed that after the unfolding of $E\varepsilon34$ TSP via urea-acid, renaturation of the protein to its native conformation was possible within few hours. This unique characteristic of refolding of $E\varepsilon34$ TSP which is similar to that of the P22 phage tailspike protein is of special interest to protein scientists and can also be exploited in antimicrobial-protein engineering.

Keywords: Renaturation; Denaturation; Phage; Fusion protein; Podoviruses

Introduction

The epsilon 34 (ε 34) phage is a dsDNA tailed-bacteriophage, carrying a 43.016Kbp in its capsid that expresses the coat protein, the tail machinery (which includes the tailspike proteins) that hydrolyzes the LPSs of its host, alongside several other structural and regulatory products. The entire sequence and the structural components of the phage have briefly been studied [1-3]. Most of these studies have been limited to sequence similarities between the P22-like phages to which ε 34 belongs [1], lysogeny and lysogenic conversion abilities of ε 34 and its phylogenetic relative known as ε 15 on their host *S. newington* [4,5]. Belonging to the temperate phages, ε 34 is known to often alter its host cell surface antigens upon lysogenization in a process called "lysogenic conversion [1,5-7]. This process is known to

block superinfection of the host by the same phage specie [8,9]. Modification of surface antigens has been demonstrated to be an important factor in bacterial virulence. For instance, Allison and Verma showed that serotype-converting bacteriophages and O-antigen modification in *Shigella flexneri* is an important virulence determinant that is conferred by temperate bacteriophages [10]. E34 phage can both undergo lytic and lysogenic pathways [1]. Exploiting phage or phage components as antimicrobial agents has received resurgence in recent times [2,11-13]. For instance, phage lysin has been employed as an antimicrobial against multidrug-resistant *Acinetobacter baumannii* [13] *Streptococcus pneumoniae* [14], biofilms of *streptococcus suis* [15]. The tailspike of

some phages too have been explored as antimicrobial agents [16-18]. In our previous communication, we demonstrated that an engineered tailspike protein from E34 phage termed as EE34 TSP showed potent antimicrobial activity against Salmonella newington, and more so, could prevent biofilm formation on decellularized tissue [2,3]. In this work, we present the unique characteristics of this protein under urea-acid treatment, a condition that recapitulates the gastrointestinal conditions as well as the excretory system [19,20] in vitro. Whiles there exist huge information on the destabilization power of urea on most protein, and while inference has been drawn based on the ability of aqueous urea to interact with both polar and nonpolar components of proteins, no data exist on the effect of urea, or urea-acid on E34 TSP. Hence, this work seeks to unravel the protein unfolding pathway via urea-acid. We further investigate the ability of E34 TSP to renature into its native state after complete denaturation via urea-acid. Similar works by Seckler and his colleagues in 1989 demonstrated that the trimeric Salmonellae phage tailspike protein, known as P22 TSP, can reconstitute in vitro after the protein had been unfolded via urea-acid [21]. Finally, we demonstrate via computational analysis the interaction between urea molecules and E34 TSP. Thus in summary, in this study, we investigated in vitro; 1) the unfolding pathway of Eɛ34 TSP under the influence of urea tinkered with acid and 2) the ability of the Eɛ34 TSP to refold back to biologically active trimeric state after it had been unfolded completely via urea-acid denaturation, and 3) finally to we reconfirm via in silico analysis the interaction between urea molecules and the hydrophobic core of Ee34 TSP.

Materials and Methods

The cloning and validation of the clone of the $E\epsilon 34$ TSP has been carried out and published in a previous communication [2].

Urea-Acid Denaturation of EE34 TSP

Samples of Eɛ34 tailspike protein in 50 mM Tris-HCl solution were titrated with urea-acid to a final concentration of 5 M urea and 2 mg/ml of the protein, and the pH adjusted to 3, 4, 5 or 6 via dropwise addition of 6N HCl. The reaction was allowed to run at room temperature and aliquots withdrawn at set time points; 0, 1, 2.5, 5, 10, 15, 20, 25, 30, and 60 minutes. Withdrawn aliquots were quickly mixed with stop buffer consisting of 50 mM Tris-HCl, 0.2% SDS, pH8 to a final concentration of 20 µg/ml. The large dilution acting as a buffer brought the acidic pH to neutrality. All samples were stored at -20°C until all experimental samples from every set time point was taken. Samples from each time point were mixed with loading buffer consisting of 50 mM Tris-HCl, 2% SDS, 5% 2-mercaptoethanol, 10% glycerol, 0.03% bromophenol blue, pH 7 and then electrophoresed in 10% polyacrylamide gel for 2 hours. Experiments were run in triplicates. Gels were stained with Coomassie blue and gel bands images acquired using Bio-Rad ChemiDoc XRS imaging system connected to PC operating Quantity One software. Densitometric values were acquired and plotted on graphs.

Renaturation of EE34 TSP

2 mg/ml of E ε 34 tsp samples was denatured in 5 M urea, 50 mM Tris (pH 3.0) for 30 minutes at 30°C. Renaturation of the TSP was commenced by initial 15-minute dialysis of the denatured protein samples in excess buffer and followed with 20-fold dilution of the unfolded TSP using the

renaturation buffer consisting of 50 mM Tris-HCl, 1mM EDTA, and pH 7.0 at $10 \pm 4^{\circ}$ C to generate a final concentration of 100 µg/ml of TSP in solution. The refolding process was allowed to proceed at 10°C with slight fluctuations in temperature at $\pm 4^{\circ}$ C, and aliquots withdrawn at specific time points and refolding reaction stopped by the addition of loading buffer consisting of 0.2 mg/ml SDS, 0.16 M Tris/HCl, 250 mg/ml glycerol, 0.1 µg/ml bromophenol blue, pH 7.0 then stored at -20°C until all samples for all specified set time points were collected. All samples were then loaded in to 10% polyacrylamide gel and electrophoresed for two hours and 30 minutes at 100 volts. Protein bands were visualized by Coomassie blue staining. Gel images were acquired using Bio-Rad ChemiDoc XRS imaging system connected to PC operating Quantity One software. Densitometric values were acquired and plotted into graphs.

Homology Modeling of ɛ34 TSP

Protein sequence extraction homology modeling of E34 TSP and urea 2D structure extraction

The protein sequences of £34 TSP (Gene ID: 7353089) was extracted from the NCBI protein database (http://www.ncbi.nlm.nih.gov/protein/). The 3D structural modeling was performed by using Swiss-Modeler (http://swissmodel.expasy.org/) an online homology modeling and model evaluation program (and work published in a previous investigation of this protein [2]. Subsequently the model's quality and validation of the model were assessed using structure assessment methods such as the QMEAN [22] and Ramachandran plot analysis (http://mordred.bioc.cam.ac.uk). The 2D structure of urea (PubChem CID: 1176) was extracted from PubChem database (https://pubchem.ncbi.nlm.nih.gov/).

Homology modeling of E34 TSP

The wild type E34 TSP sequence was extracted from NCBI website and modeled following a similar process by Filiz et al. [23]. In short, the E34 TSP sequence was blasted against PDB to extract homologous templates for the modeling. The bifunctional P22 TSP, a homotrimer with PDB ID 2XC1.1.A [24] possessing 69.83% amino acid identities was used to model the HBD of the E34 TSP. The modeling of HBD of E34 TSP its 3D structure was carried out using the Swiss-Modeler (http://swiss-model.expasy.org/) program [23,25,26]. The quality of the HBD model was evaluated to show a QMEAN of 0.66 ± 0.05 and a Ramachandran plot analysis as shown in Figure pred2B, the molProbity analysis showed a clash score of 3.09, a 96.83% Ramachandran favored structure, 0.00% Ramachandran outliers, 1.66% rotamer outliers and 0 bad bonds out of 2842 bonds. For the local quality estimate, the first 12 amino acids showed QMEAN of less than 0.3, whereas the rest of the structure showed amino acids having a local QMEAN greater than 0.4.

Molecular Docking analysis of E34 TSP- urea

To investigate the interactions between the urea (PubChem CID: 1176) and £34 TSP, the 2D structure of urea was downloaded from the PubChem database and docked to £34 TSP using PyRx (an open-source software for performing virtual screening that combines AutoDock Vina, AutoDock 4.2, Mayavi, Open Babel etc). The modeled structure of £34 served as the

receptor. The modeled E34 TSP was prepared using AutoDock Vina wizard, whereas the urea molecule was prepared by using the Open Babel tool. In summary, the 2D urea molecular structure was minimized and converted to a pdbqt format before uploading as a ligand. In the case of the receptor, the bond orders were assigned, and charged hydrogen atoms added to the protein. The receptor structure was also minimized using the AutoDock Vina wizard. Additionally, proteins were loaded into PyRx and converted to receptors, the receptor grid boxes were generated in PyRx using the build-in Vina Wizard module, the grid box was maximized to cover all active sites of the E34 TSP head binding domain (HBD) (Which demonstrate a globular structure and shows high hydrophobic regions). AutoDocking of urea to the HBD of E34 TSP was made by using the AutoDock wizard in-built in PyRx program to blind dock the urea molecule to the HBD of the E34 TSP with an exhaustiveness of 9. The free energies generated in docking results were saved and the models were generated were exported to Biovia Discovery Studio software (version; 21.1.0.278) for specific atomic-atomic interaction analysis between the ligand and the receptor.

were evaluated using P-values of Student's t-test (one-tailed, two samples of unequal variance, significance level α =0.05 was used unless otherwise stated).

Results

Urea Denaturation Profile

As can be observed Figure 1A, and Figure 1B, denaturation of $E\epsilon 34$ TSP occurred very quickly, with protein species existing predominantly in monomeric forms in the 5th minute onwards. A faint band can however be observed in the 2.5-minute mark for the 5 M, pH 4 (Figure 1B), but absent in the 5 M, pH 3, an indication of an incomplete unfolding of all species at the 5 M, pH4 in this time point, but a complete denaturation of all samples at the 5 M, pH3 at the same time point. The higher pH 5 and pH 6 as depicted by Figure 1C and Figure 1D show all samples in their trimeric states, indicative of resistance to denaturation at these conditions for all time points.

Statistical analyses

For all statistical data, values were derived from multiple measurements (from replicates of 3 experiments) and averaged, the standard deviations To understand in detail, the denaturation characteristics of $E\epsilon 34$ TSP under these set conditions, we repeated this experiment three times and band densities were quantitatively analyzed.



stock urea to urea set concentration of 5M, and the pH tinkered to 3, 4, 5 or 6 using 6NHCl. At the set time points, aliquots were withdrawn and quickly mixed with stop buffer consisting of 50 mM Tris-HCl, 0.2% SDS, pH8 to a final concentration of 20 μ g/ml and stored temporarily at -20 °C until all sample reactions were quenched and collected. Samples were then electrophoresed via an SDS-polyacrylamide gel and imaged using the ChemiDoc XRS installed with Quantity one software. UHUT = Unheated untreated Ee34 TSP samples. HUT = Heat but untreated Ee34 TSP samples. PPPS = Pre-stained Perfect Protein Standard. 0, 2.5, 5, 10, 20, 30, 40, and 60 are time recorded in minutes.

At pH 3, in 5 M urea (Figure 2), the quickest and sharpest drop of TSP species from trimeric forms to monomers is recorded. At an initial high of over 80% trimers, 60 seconds of treatment under this condition pulled the trimeric species curve down to almost 0%. No record of trimeric forms is seen with passing time afterward. While a quick drop in trimeric forms occurred in a space of a minute, this same space of time gave a sharp rise in the monomers, registering an average of 100% for monomers in only 1 min

ute of treatment. This percentage is maintained for monomers for all other time points extended from the 1-minute mark. The time point at which the two curves; monomeric species curve and trimeric species curve, intersected was recorded to be 30 seconds and registered a 50% split for both species type. This is indicative of complete denaturation at the highest speed for our TSP compared to all other conditions tested against the protein in this study.



Figure 2. Chart following the kinetics of denaturation of $E\epsilon 34$ TSP in 5 M urea, pH 3. Chart following the kinetics of denaturation of $E\epsilon 34$ TSP in 5M urea, pH 3 compiled from the densitometric values obtained from bands of triplicate experiments. %TRM D = percentage trimeric $E\epsilon 34$ TSP species under denaturation condition of urea-acid, %MM D= percentage $E\epsilon 34$ TSP monomers under denaturation condition of urea-acid. p-value = 0.05, n= 3.

As depicted in Figure 3, at room temperature $E\epsilon 34$ TSP trimeric form existed in equilibrium with their monomeric counterparts, holding a huge share of approximately 90% while 10% represented monomers. However, in less than a minute, treatment of $E\epsilon 34$ TSP to 5 M urea, pH 4 produced a drop in the percentage trimers to approximately 77.5%, at 2.5 minutes less than 35% TSP exist in trimeric state. The rest of the time points maintained an average of approximately 34% trimeric state TSP species. Monomeric species how

ever stood at 12.5% at 0-time point, increased steeply to 65.1% in only 2.5 minutes and stayed approximately the same for the entire experimental period of 120 minutes. The inverse relationship between the monomeric forms and the trimers are best seen at the early seconds of the treatment, with these two curves intersecting at the 89th second to record a percentage of 50% trimers and 50% monomers.





Treatment of E ε 34 TSP to 5M urea, pH 5 (Figure 4) produced a simple almost linear curve with nearly a zero slope. In this study, at 0-time point, the E ε 34 TSP samples used had 70.5% trimeric and 29.5% monomeric species, then within half a minute, treatment of E ε 34 TSP with 5M urea, pH 5 resulted in a slight increase in percentage of trimeric TSP population, then a 2.5-minute treatment showed an increased to about 87.5% trimeric TSP, 5 minutes gave an average percentage of 92.3% and a 94.9% at 10 minutes. Over an average of 99.3% trimers were recorded at time points exceeding the 10-minute mark. The monomeric forms however, decreased with time, starting slightly over 29.4% and dropping continuously to a low of 5% at 10 minutes, and 0.7% at the 30th minute. It maintained an average low of 2.6% at the 60th minute mark.

This seems to indicate that at pH 5, in 5M urea, Eɛ34 TSP is resistant to denaturation, although it also seems to show a pattern in which some species of monomers are forced into association to form trimers.



As shown in Figure 5, the trimeric species indicated 87.5% abundance at 0-time point, an incremental drop to an average of 84.8% was recorded at just 10 seconds and at 2.5 minutes later 84.9% abundance of the trimers was registered. A slight gain in percentage to 88.9% came at the 5th minute, and average abundance for the rest of the time points stayed at an average of 88% beyond the 5th minute mark.

The monomers at the other hand registered an average of 12.5% at the

0-time point at room temperature and peaked at 15.1% at 2.5 minutes then

dropped down to 11.1% at the 5-minute and continue to exist in the lower percentages below 10% average for all readings beyond the 5-minute mark.

This demonstrates that EE34 TSP is very stable in 5M urea, pH 6. Repeating the same experimental conditions with an extended period (from 60 minutes to 180 minutes) was carried out to gain insight to the nature of TSP at 5M urea, pH 6 for extended time points (Data not shown). After 180 minutes, at least 97.9% of TSP existed in their trimeric form in all samples while less than 2.1% remained as monomers.



To appreciate the kinetics and gain greater insight of the characteristics of the protein under the varying conditions for the study, a graph focusing in the first five minutes of the kinetics are made for comparison, thus providing for a holistic view comparing the kinetics of the four denaturation curves (Figure 6).

5 and 6 respectively produced less than 20% monomeric species of $E\epsilon 34$ TSP on the average. An inverse relationship existing between monomeric species and the trimers saw the sharpest fall of trimers in the 5 M Urea, pH3, followed by 5 M Urea, pH4. The 5 M Urea, pH5 and pH6 maintained an almost flat curve, registering a high of over 80% trimers in average.

As shown in Figure 6, 5 M Urea, pH 3 gave the sharpest gradient in monomeric species production followed by 5 M Urea, pH4. 5 M Urea at pH



Renaturation Kinetics of EE34 TSP

Protein folding, unfolding, tagging, proteolysis, aggregation and reconstitution are simultaneous events occurring in the cytosolic milieu of cells [21,27,28]. For instance, while aggregation could arise due to hydrophobic interactions between "wrongly ex- posed" protein surface areas produced by temperature and ionic strength [29] *in vivo*, the cell employs several biochemical processes to rescue protein from aggregating [30], more so, *in vitro* folding experiments have demonstrated that cytosolic globular proteins possess small folding equilibrium constants and do unfold and refold many times during their life cycle [21]. The reconstitution of the P22 tailspike protein from denatured chains *in vitro* has been documented [21].

To investigate the refolding pathway of extended ε 34 TSP, this protein samples were subjected to complete denaturation via 5M urea, pH 3 for half an hour and subjected to dialysis in excess buffer consisting of 50 mM Tris, 1 mM EDTA, pH 7.0 at room temperature for a brief period of 15 minutes, afterward; samples were 20-fold diluted by the renaturing buffer (50 mM Tris, 1 mM EDTA, and pH 7.0 pre-chilled to 10°C) to commence renaturation. Reconstitution reaction was incubated at 10±4°C. Renaturation was stopped at the following designated time points by the addition of a quenching solution consisting of 0.2 mg/ml of SDS, 50 mM Tris/HCl, 250 mg/mL glycerol, 0.1 µg/ml bromophenol blue, pH 7; The set time points were 0-minute, 10 minutes, 30 minutes, 60 minutes, and 3 hours, 4 hours, 5 hours, and 9 hours and results of each of these time points shown in lane 2 to lane 10 respectively. Lane 1 is the standard protein ladder (Colored Prestained Protein Standard (CPPS), New England Biolabs). The samples were separated by electrophoresis in 10% polyacrylamide gel and visualized by Coomassie blue staining.

A similar method has been employed by Seckler and co. in their renaturation study of P22 [21], the exception however in this case was the additional use of dialysis method to precede the commencement of the renaturation process instead of direct initial 50-fold dilution employed by the said investigators. Dialysis tubes served three major advantages; 1) There is no loss of protein concentration during the dialysis with the concomitant quick reduction in the denaturant (urea) concentration. 2) Protein samples were incubated in an excess of 5M urea, pH 3 which served to quickly denature the protein within half an hour, the high acidity of the solution was quickly buffered to neutrality during the brief period of dialysis. Previous denaturation study showed that protein samples were completely denatured in this process within the first 2 minutes; 3) Simple protein detection/visualization method such as Coomassie blue staining can be employed instead of the costlier silver staining process. The renatured samples were run in 10% polyacrylamide gel. The untreated control as in lane 2 Figure 6 migrated to the position characteristic of the Eɛ34 TSP at its trimeric state.

SDS-Page Analysis of Renaturation



Figure 7. The renaturation kinetics of ϵ 34 TSP into its trimeric state as observed in a 10% SDS-PAGE from fully denatured samples. Samples of EE34 TSP were denatured in 5M urea, pH 3 for half an hour, the renaturation process was commenced via first 15 minutes' dialysis and followed with 20-fold dilution of Ee34 TSP samples using renaturation buffer consisting of 50 mM Tris, 1 mM EDTA, and pH 8.0 at 10 ±4°C. Samples were withdrawn from the renaturation reaction at set time points and the reaction quenched by the addition of quenching solution composed of 50 mg/ml SDS, 50 mM Tris-HCl, 250 mg/ml glycerol, 0.1 mg/ml bromophenol blue, pH 7.0. The set time points were 0 minute, 2.5 minutes, 5 minutes, and 1 hour, 2 hours, 5 hours, and 9 hours as shown in lane 3, lane 4, lane 5, lane 6, lane 7, lane 8, lane 9 and lane 10 respectively. Lane 1 represented Prestained Perfect Protein Standard (PPPS) (Novagen), lane 2, is unheated and untreated control sample. Samples were run in 10% SDS-PAGE, visualized by Coomassie blue R-250 staining. Gels were scanned using Chemi-Doc fitted with Quantity One software.

Quantitative Analysis of Renaturation of $E\epsilon 34$ TSP after Samples

To quantitatively analyze the renaturation of EE34 TSP after the protein samples had been subjected to an initial 5 M urea, pH 3 denaturation processes for half an hour we followed EE34 TSP refolded species using SDS-PAGE and quantitated the densitometric values of the arising bands. As shown in Figure 8, the renaturation characteristic is indicative of hysteresis; a quicker denaturation process that occurred in less than 60 seconds in 5M urea, pH 3, took over 300 minutes to produce 80.6% trimers again, and about 275 minutes of renaturation time to produce 50% native conformers. While in less than 30 seconds, $E\epsilon34$ TSP recorded 50% trimers in the denaturation process. It was also observed that 100% trimeric forms were

not possible within the reach of our experimental time frame even after we allowed renaturation to continue for an extended period. Upward inflection of our trimeric species curve started at the 180-minute mark, recording a slight increase in trimeric species from approximately 0.2% to 1.3%, and then rose steadily to 26.2% average at the close of the 240th minute. The 300-minute mark recorded 80.6% trimeric species. The downward trajectory of the monomeric species also started at the 180th minute mark, registering a lowered percentage of 98.6%, a drop from 99.8%. Then 73.8% monomers were recorded at the close of the 240th minute and a lowest of 19.4% at the end of the 300-minute time point.



Figure 8. Renaturation kinetics of $E\epsilon_34$ TSP after complete urea-acid denaturation. Renaturation profile for $E\epsilon_34$ TSP after samples had been subjected to an initial 5 M urea, pH 3 denaturation processes for half an hour. Renaturation process was commenced via 20-fold dilution of $E\epsilon_34$ TSP samples using renaturation buffer consisting of 50 mM Tris, 1 mM EDTA, and pH 8.0 at 10 ±4°C. Samples were withdrawn from the renaturation reaction at set time points and the reaction quenched by the addition of quenching solution composed of 50 mg/ml SDS, 50 mM Tris-HCl, 250 mg/ml glycerol, 0.1 mg/ml bromophenol blue, pH 7.0. The set time points were 0 minute, 10 minutes, 30 minutes, 60 minutes, and 3 hours, 4 hours, 5 hours. Data were obtained via averaging 3 replicates of densitometric values of electrophoretic bands after SDS-PAGE electrophoresis of renaturation samples arising from three separate experiments.

Computation Analysis of Urea- ɛ34 TSP Interaction

Calorimetric titration analysis of the interaction of urea with proteins has been studied and demonstrates that urea binds to proteins leading to their unfolding. It has been revealed that the binding of urea by protein follows a drop in enthalpy and entropy [31]. Analysis of the binding sites of urea to the studied proteins showed that urea binds by forming hydrogen bonding [31] which significantly restricts the conformational freedom within the polypeptide chain. In a molecular dynamics simulations of chymotrypsin inhibitor in 8 M urea at 60°C the protein unfolded; however it retained its native structure in water at the same temperature [32]. They also observed that the overall process of unfolding in urea resembles the protein's unfolding process in thermal denaturation simulations above the protein's Tm of 75°C [32]. Notably, they demonstrated that the initial unfolding step was the expansion of the hydrophobic core [32] which led to the solvation of the core by water and urea.

In this work, as shown in Figures 10 and 11, urea binds to the hydrophobic core of the HBD of ε 34 TSP via the amino acid residues as depicted in Table 1. It is clear from Table 1, that urea is engaged in hydrogen bonding with the protein. The *in vitro* characterization of the denaturation of E ε 34 TSP by urea alone showed that urea could not denature E ε 34 TSP at room temperature, even at higher concentrations up to 7 M (data not shown). However, at lower pH conditions, urea at 5 M was able to denature E ε 34 TSP (Figures 1A, 1B, 2 and 3). Proving that urea might require an initial expansion of the hydrophobic core of E ε 34 TSP similar to what has been described by Bennion and Daggett [32], in order to access and bind to the shielded hydrophobic core.



Figure 9. Putative head binding domain (HBD) of £34 TSP (grey) docked to urea molecules (red). (A) Cartoon structure of the HBD of putative £34 TSP interacting with urea in red. (B) Space filled structure of the HBD of £34 TSP showing urea interacting in the hydrophobic core of the protein. (C) An enlarged view of the urea enclosed by the hydrophobic interacting atoms of HBD of £34 TSP.



Figure 10. Putative head binding domain (HBD) of £34 TSP (grey) docked to urea molecules (red). (A) Cartoon structure of the HBD of putative £34 TSP interacting with urea in red enclosed in a hydrophobic binding site. (B) The interaction of urea with the specific interaction atoms of the HBD of £34 TSP, also depicting the degree of hydrophobicity of the binding cavity. (C) 2D depiction of urea molecule binding to four amino acids in the highly hydrophobic binding pocket of HBD of £34 TSP. The free relative energy recorded was -3.4 kcal/mol. Specific interactions are shown in Table 1.



Figure 11. Putative head binding domain (HBD) of £34 TSP (grey) docked to urea molecules (red). (A) Cartoon structure of the HBD of putative £34 TSP interacting with urea in red at a slightly less hydrophobic region compared to Figure 10. (B) The interaction of urea with the specific interaction atoms of the HBD of £34 TSP, also depicting the degree of hydrophobicity of the binding cavity. The free relative energy recorded was -2.9 kcal/mol. Specific interactions are shown in Table 1.

Model	Residue	Bond distance/Å	Bond type	∆G/kcal/mol
1	GLN15	1.98	Hydrogen Bond	-3.4
1	ALA80	2.83	Hydrogen Bond	
1	SER14	2.19	Hydrogen Bond	
1	ALA80	2.73	Hydrogen Bond	
1	PHE82	1.88	Unfavorable bond	
6	GLN86	1.85	Hydrogen Bond	-2.9
6	PRO48	2.52	Hydrogen Bond	
6	THR84	2.42	Hydrogen Bond	
6	VAL85	1.69	Unfavorable bond	

Table 1. The site-specific interaction of urea with the HDB of £34 TSP, depicting the interacting amino acids, the bond distances, the bond types recorded as well as the relative free energies predicted to be released upon interaction of urea and £34 TSP.

Discussion

The analysis of the urea-acid denaturation process of $E\epsilon 34$ TSP have helped identify two major aspects of the protein; 1) first it proofs that, the protein have no intermediate since no intermediary band was noticeable, however two forms of the protein were recorded, thus the monomeric and the trimeric states of the protein. 2) It showed that the unfolding and reconstitution follows separate unique pathways. The renaturation studies demonstrated that $E\epsilon 34$ TSP can be renatured with a substantial good yield under refolding conditions *in vitro*. While it took 48 hours for P22 TSP to achieve 80% renaturation [21], 5 hours of renaturation of $E\epsilon 34$ TSP produced similar results. In the case of P22 TSP renaturation began slowly at 3-hour time point, and progressively increased with time to achieve complete refolding at 96 hours [21], $E\epsilon 34$ TSP however did not achieve 100% renaturation even after extended time points, however, began similarly at the 3-hour time mark, and progressed quickly to achieve 80.6% trimeric forms at 5-hour time point. In the limits of experimental errors, and the slight difference in experimental conditions for which our protein was subjected to e.g. temperature fluctuation within the limits of $10 \pm 4^{\circ}$ C and a lower dilution of 20-fold, whereas the study by Seckler and his colleagues on P22 TSP received a constant temperature of 10° C and with higher dilution of 50fold, it seems plausible to assume that Eɛ34 TSP refolds quicker than P22 TSP. An argument that our lower dilutions (20-fold instead of the 50-fold dilution used by Chen and King) might have resulted in higher concentration of our polypeptide chains, which might present higher frequency of collision and association into fully matured trimeric species, whereas the higher dilution of P22 TSP decreased the polypeptide collision frequency and hence their frequency of association and oligomerization is feasible. Nonetheless, it is interesting to observe that both studies recorded an initial reconstitution starting at the 3-hour time point. The reconstituted products of EE34 TSP after the 300-minute mark and the 540-mark showed nativelike characteristics since these species were resistant to SDS under room temperature and hence migrated at a trimeric level as demonstrated via the SDS-PAGE result (Figure 7).

This in vitro renaturation system would enable a deeper understanding of the folding and association pathway of the EE34 TSP. Analytic comparison of the in vitro renaturation of both P22 TSP and EE34 TSP further elucidates the functional and structural similarities these two proteins share. A major drawback of this study is the fact that this is conducted in an in vitro condition, thus will substantially differ from an in vivo folding and chain association processes since in in vivo conditions, nascent polypeptides are assisted to fold at unfavorable conditions by molecular chaperone [33]. The obvious question that lingers here is, to what extend will such information regarding in vitro refolding process guide and inform our understanding of the complex in vivo folding process in the ɛ34-gp19 translational process in the life cycle of the ɛ34 phage? Since there exist a complex and dynamic cellular environment of the host cell compared to the test tubes, and the presence of molecular chaperones in vivo to assist unfolded TSPs or nascent polypeptides fold into their proper conformations but absent in vitro further compounds issues. The chances of post proteolytic processing of the TSPs if administer in vivo are not captured in these in vitro assays, and these can possibly add up to confound and or limit our full grasp of the true nature and property of folding, unfolding, and refolding of our TSP in vivo when administered orally or intravenously as an antimicrobial. However, it can be glimpsed from this study that even within the limits of in vitro experimentation, reconstitution of Ee34 TSP is possible.

The *in-silico* studies (Figures 9, 10 and 11) further support established knowledge that urea binds to solvent inaccessible hydrophobic regions, thus destabilizing the protein's core and subsequently leading to the protein's denaturation. The EE34 TSP however showed high resistance to urea denaturation at higher pHs (Figures 1, 4 and 5).

Conclusion

In conclusion, this work unraveled the $E\epsilon 34$ TSP unfolding, and renaturation pathways, demonstrating the high stability of the protein against urea denaturation. Furthermore, this work demonstrated the ability of $\epsilon 34$ TSP to renature into its native SDS resistant state after complete denaturation via urea-acid. Finally, we demonstrated via computational analysis the interaction between urea molecules and $\epsilon 34$ TSP. Thus in summary, in this study, we investigated *in vitro*; 1) the unfolding pathway of $\epsilon 34$ TSP under the influence of urea tinkered with acid and 2) the ability of the $\epsilon 34$ TSP to refold back to biologically active trimeric state after it had been unfolded completely via urea-acid denaturation, and 3) finally to we reconfirm via *in silico* analysis the interaction between urea molecules and the hydrophobic core of $\epsilon 34$ TSP. These discoveries taken together shows that $\epsilon 34$ TSP is a stable protein and shows similar characteristics as the well-studied P22 TSP. Thus, providing evidence that the use of this protein as an antimicrobial agent against *S. newington* in an *in vivo* system could be feasible.

Acknowledgements

We thank the department of Microbiology, Faculty of Science, Mathematics and Technology, Alabama State University for timely support.

Conflict of Interest: All authors declare that they have no conflicts of interest.

Ethical Approval: This article does not contain any studies with human participants or animals performed by any of the authors.

References

- Uetake H, Luria SE, Burrous JW. Conversion of somatic antigens in Salmonella by phage infection leading to lysis or lysogeny. Virology. 1958, 5: 68-91.
- Ayariga JA, Gildea L, Villafane R. E34 Phage Tailspike Protein is Resistant to Trypsin and Inhibits and Salmonella Biofilm Formation. Preprints. 2021.
- Ayariga JA, Gildea L, Wu H, Villafane R. The £34 Phage Tailspike Protein: An *in vitro* characterization, Structure Prediction, Potential Interaction with *S. newington* LPS and Cytotoxicity Assessment to Animal Cell Line. bioRxiv. 2021.
- Villafane R, Zayas M, Gilcrease EB, Kropinski AM, Casjens SR. Genomic analysis of bacteriophage epsilon 34 of Salmonella enterica serovar Anatum (15+). BMC Microbiol. 2008, 8: 227.
- Uetake H, Nakagawa T, Akiba T. The relationship of bacteriophage to antigenic changes in Group E salmonellas. J Bacteriol. 1955, 69: 571-579.
- Uetake H. The genetic control of inducibility in lysogenic bacteria. Virology. 1959, 7: 253-262.
- Uetake H, Hagiwara S. Genetic cooperation between unrelated phages. Virology. 1961, 13: 500-506.
- Wright A, Barzilai N. Isolation and haracterization nonconverting mutants of bacteriophage epsilon 34. J Bacteriol. 1971, 105: 937-939.
- Wright A. Mechanism of conversion of Salmonella O antigen by bacteriophage epsilon 34. J Bacteriol. 1971, 105: 927-936.
- Allison GE, Verma NK. Serotype-converting bacteriophages and O-antigen modification in *Shigella flexneri*. Trends Microbiol. 2000, 8: 17-23.
- Moradpour Z, Ghasemian A. Modified phages: novel antimicrobial agents to combat infectious diseases. Biotechnol Adv. 2011, 29: 732-738.
- Fenton M, Ross RP, McAuliffe O, O'Mahony J, Coffey A. Characterization of the staphylococcal bacteriophage lysin CHAPK. J Appl Microbiol. 2011, 111: 1025-1035.
- Thandar M, Lood R, Winer BY, Deutsch DR, Euler CW, Fischetti VA. Novel engineered peptides of a phage lysin as effective antimicrobials against multidrug-resistant *Acinetobacter baumannii*. Antimicrob Agents Chemother. 2016, 60: 2671-2679.
- Resch G, Moreillon P, Fischetti VA. A stable phage lysin (Cpl-1) dimer with increased antipneumo-coccal activity and decreased plasma clearance. Int J Antimicrob Agents. 2011, 38: 516-521.
- Meng X, Shi Y, Ji W, Meng X, Zhang J, Wang H, et al. Application of a bacteriophage lysin to disrupt biofilms formed by the animal pathogen *Streptococcus suis*. Appl Environ Microbiol. 2011, 77: 8272-8279.
- Olszak T, Shneider MM, Latka A, Maciejewska B, Browning C, Sycheva LV, et al. The O-specific polysaccharide lyase from the phage LKA1 tailspike reduces Pseudomonas virulence. Sci Rep. 2017, 7: 1-14.

- Waseh S, Hanifi-Moghaddam P, Coleman R, Masotti M, Ryan S, Foss M. et al. Orally administered P22 phage tailspike protein reduces Salmonella coloni-zation in chickens: prospects of a novel therapy against bacterial infections. PLoS One. 2010, 5: e13904.
- Lee IM, Tu IF, Yang FL, Wu SH. Bacteriophage Tail-Spike Proteins Enable Detection of Pseudamin-ic-Acid-Coated Pathogenic Bacteria and Guide the Development of Antiglycan Antibodies with Cross-Species Antibacterial Activity. J Am Chem Soc. 2020, 142: 19446-19450.
- Miller TL, Wolin MJ. Fermentation by the human large intestine microbial community in an *in vitro* semicontinuous culture system. Appl Environ Microbiol. 1981, 42: 400-407.
- 20. Lu HY, Ning XY, Chen YQ, Han SJ, Chi P, Zhu SN, et al. Predictive value of serum creatinine, blood urea nitrogen, uric acid, and β2-microglobulin in the evaluation of acute kidney injury after orthotopic liver transplantation. Chin Med J (Engl). 2018, 131: 1059-1066.
- Seckler R, Fuchs A, King J, Jaenicke R. Reconstitution of the thermostable trimeric phage P22 tailspike pro-tein from denatured chains *in vitro*. J Biol Chem. 1989, 264: 11750-11753.
- Benkert P, Biasini M, Schwede T. Toward the estimation of the absolute quality of individual protein structure models. Bioinformatics. 2011, 27: 343-350.
- Ertugrul F, Koç I. *In silico* sequence analysis and homology modeling of predicted beta-amylase 7-like protein in Brachypodium distachyon. LJ BioSci Biotechnol. 2014, 3: 61-67.
- Seul A, Muller JJ, Andres D, Stettner E, Heinemann U, Seckler R. Bacteriophage P22 tailspike: structure of the complete protein and function of the interdomain linker. Acta Crystallographica Section D: Biological Crystallography. 2014, 70: 1336-1345.
- Arnold K, Bordoli L, Kopp J, Schwede T. The SWISS-MODEL workspace: a web-based environment for pro-tein structure homology modeling. Bioinformatics. 2006, 2: 195-201.

- Bordoli L, Kiefer F, Arnold K, Benkert P, Battey J, Schwede T. Protein structure homology modeling using SWISS-MODEL workspace. Nat Protocols. 2009, 1: 1-13.
- Frydman J. Folding of newly translated proteins *in vivo*: the role of molecular chaperones. Annu Rev Biochem. 2001, 70: 603-647.
- Clark PL, King J. A newly synthesized, ribosome-bound polypeptide chain adopts conformations dissimilar from early *in vitro* refolding intermediates. J Biol Chem. 2001, 276: 25411-25420.
- van der Kamp MW, Daggett V. Pathogenic mutations in the hydrophobic core of the human prion protein can promote structural instability and misfolding. J Mol Biol. 2010, 404: 732-748.
- Soto C, Pritzkow S. Protein misfolding, aggregation, and conformational strains in neurodegenerative diseases. Nat Neurosci. 2018, 21: 1332-1340.
- Makhatadze GI, Privalov PL. Protein interactions with urea and guanidinium chloride: a calorimetric study. J Mol Biol. 1992, 226: 491-505.
- Bennion BJ, Daggett V. The molecular basis for the chemical denaturation of proteins by urea. PNAS. 2003, 100: 5142-5147.
- Eustace BK, Jay DG. Extracellular roles for the molecular chaperone, hsp90. Cell cycle. 2004, 3: 1096-1098.
- Zou Q, Habermann-Rottinghaus SM, Murphy KP. Urea effects on protein stability: hydrogen bonding and the hydrophobic effect. Proteins. 1998, 31: 107-115.

Submit your manuscript at http://enlivenarchive.org/submit-manuscript.php

New initiative of Enliven Archive

Apart from providing HTML, PDF versions; we also provide video version and deposit the videos in about 15 freely accessible social network sites that promote videos which in turn will aid in rapid circulation of articles published with us.