

Gerontology and Nursing in the Emergency Department

Gerard Charles, RN, MSc^{1,2*}, Sophia Pei Wen Lau, RN, MSc^{1,2}, and Muhammad Saifuddin Bin Supandi, RN, BN²

¹S R Nathan School of Human Development, Singapore University of Social Sciences

²Singapore General Hospital, Department of Emergency Medicine

***Corresponding author:** Gerard Charles, S R Nathan School of Human Development, Singapore University of Social Sciences, Singapore General Hospital, Department of Emergency Medicine, E-mail: gerard.charlesix@gmail.com

ORCID: <https://orcid.org/0000-0003-2545-292X>

Received Date: 29th May 2021

Accepted Date: 18th June 2021

Published Date: 25th June 2021

Citation: Charles G, Lau PWS, Saifuddin MS. (2021) Gerontology and Nursing in the Emergency Department. Enliven: J Anesthesiol Crit Care Med 8(4): 005.

Copyright: © 2021 Gerard Charles. This is an Open Access article published and distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Abstract

Background: With a rapid rise in our older adult population globally and due to their multimorbidities, our older adults are more likely to engage in the services provided in the emergency department at a higher rate than younger adults. The current emergency service delivery model may be ineffective against such an ageing phenomenon. Research reports older adults having one or more co-morbidities including functional decline, dementia, and frailty. Studies have shown that older adults have been undertriaged with physicians not being able to comprehend their complex needs related to their presenting complaints in the emergency department. Geriatric emergency departments have incepted worldwide to better manage this care deficiency in the rapidly ageing society around the world.

Data Sources: A search of published literature from 2010-2020 using keywords as described below was undertaken of which, relevant literature were selected for an informed review.

Implications for Nursing: Understanding geriatric emergencies can enable healthcare workers to reduce undertriaging and provide appropriate care that improves patient's health outcomes currently and in the future in the emergency department. Further education in gerontology can also be a platform for our nurses to enhance their care and thought process, likewise upskilling themselves for the future geriatric population seeking treatment. Older healthcare workers will also be able to enhance their current job scope before retirement. In-house teachings from trained gerontologists or certified programs can shed light on the special care needs of our senior citizens globally.

Conclusion: With our rapidly increasing ageing population, we can expect an influx of our older patients both from home and long-term care facilities to present to the emergency department with a wide range of geriatric emergencies. By being able to create a geriatric screening process and tailored care models, healthcare workers will be able to understand their care process and in turn, improve patients' health outcomes and provide a quicker transition of care.

Keywords: Nursing; Emergency department; Gerontology; Geriatric nursing; Geriatric emergency; Emergency nursing; Geriatric emergency unit

Introduction

Globally, the percentage of our older adults had seen a rapid rise of 9.2% in 1990 to 11.7% in 2013, with projections anticipating a further increase to 21.1% which would constitute more than 2 billion people by 2050 [1]. Throughout the past 10 years, the increase of our older adults have posed a significant burden to healthcare delivery and finances as more than 50% of older adults aged 65 years old and above have at least two chronic conditions (multi-morbidity) [2,3]. Older adults are more likely to engage in the

services provided in the emergency department (ED) at a higher rate than their younger counterparts [4]. Older adults might also have difficulties in articulating their concerns to the healthcare workers in the ED, thus, creating multiple opinions in their treatment process. This could lead to undertriage, in which delays, could pose a severe risk to the older adult's health outcomes, given the long waiting times present in EDs worldwide.

Apart from the evident chronic diseases, multi-drug regimen, and abnormal presentations, the seniors visiting the ED often have underlying issues not identifiable to the physician during examination [5-7]. Functional decline and impaired cognition, for instance, delirium or dementia, chronic diseases, frailty, falls, and polypharmacy have consequential detrimental impacts on an older adult's psychosocial and physical well-being [8-14]. In geriatric emergency patients, the risk of adverse outcomes such as hospital (re) admission, institutionalisation, and mortality are increased compared to younger patients [6,15]. This paper aimed to present a discussion of special care needs and geriatric emergencies for older adults seeking treatment in the emergency departments.

Background

What are the Special Care Needs of Older Adults?

Functional Decline:

One of the consequences of ageing is functional decline. It can be defined as the onset of new disabilities [16]. This can significantly impact an older adult's activities of daily living (ADL) or instrumental activities of daily living (IADL). Katz et al. [17] categorised six domains as measures of our ADL; bathing, dressing, toileting, transferring, continence and feeding. Lawton & Brody [18] provided a more in-depth description of our IADLs such as using the telephone, shopping, preparing food, housekeeping, doing laundry, using transportation, handling medications, and handling finances. A loss of independence in ADLs or IADLs signifies a functional decline in an individual which may either be temporary or irreversible.

Functional decline can occur through various facets, particularly during and after hospitalization. Functional decline post-hospitalization is a prevalent adverse outcome that occurs in the elderly, mainly caused by immobilization, polypharmacy, isolation, delirium, and pressure sores [19]. As informed by Covinsky et al. [20] an estimate of a third of older adults encounter functional incapacity post hospitalization. The association of acute complaints and geriatric disorders are significant indicators of adverse outcomes during hospital admission which are also often correlated with functional and cognitive impairment, institutionalisation, and mortality after discharge [21-26].

During a hospital stay deconditioning resulting in functional decline is a major problem [27]. When functional decline occurs, independence is affected in physical, cognitive, and social/emotional domains. Despite the ability to walk, older people spend most of their time in hospital, in bed [28]. Abilities for self-care when lost, are not easily regained by an older person [29-32].

Frailty

While there is no standard definition of frailty, it is generally explained as muscle weakness, fatigue, slowness, low physical activity, and unintended weight loss [33-36,130]. Gobbens et al. [37,38] also defined frailty as a dynamic state affecting an individual who experiences losses in one or more domains of human functioning. Frail older patients have higher ED conversion rates [39] that result in longer admissions [40] with a greater likelihood of re-admission [41] and higher inpatient mortality [40]. By identifying patients who are frail at the EDs, potential interventions could be initiated to ensure their safety, preventing adverse events that could potentially occur in the ED such as falls, skin tears, pressure sores, and

delirium. Interventions such as placing the patient on a trolley bed instead of a wheelchair would greatly aid in the prevention of wandering that could occur amidst the overcrowded ED.

Falls

Falls are a common reason for older adults attending the ED, and older adults with a fall-related ED visit have increased risk for subsequent falls, hospital re-admission, functional decline, and mortality [14,42-45]. Research has shown, patients who have sustained a fall are most likely to fall again. Recurrent falls are often sub-sequential falls that are not investigated the first time it occurs [46]. 30% of fall victims are predisposed to recurrent falls. Recurrent falls are described as two or more falls per annum with low attribution to a 'mechanical' (i.e. accidental) cause. Common precipitating factors for falls are attributed to issues such as co-morbidities that could be preventable. Associating factors with fear of fall 'post-fall anxiety syndrome' can occur with a near fall incident that results in self-imposed reduced mobility causing anxiety and eventually despair. These are often witnessed in seniors living alone with impaired cognition and mobility and poor gait which precipitates subsequent falls [47].

There are several factors toward fall prevention that can be modified amongst our seniors that includes poor and/or impaired mobility due to joint or medical diseases (e.g., Arthritis, Parkinson's disease, and cerebrovascular accident), polypharmacy (e.g., sedatives and antipsychotics), cognition or vision deficit (e.g., Alzheimer's or cataracts), non-elder friendly environment and improper foot gear [131]. Usually, ED assessments for falls require a targeted review to rule out acute trauma and therefore do not address modifiable risk factors to prevent recurrent falls [11,48,49]. Preventing subsequent falls is critical to stop the cascade of functional decline, loss of independence, hospitalization, and death, which frequently follows a fall. Thus, a patient with a fall that can be discharged to home should undergo a teaching session led by a geriatric nurse, gerontologist, or in some places, patient navigators. Education related to the prevention of subsequent falls should be well informed to the patient.

Dementia

Dementia originates from the Latin term 'de mens' [50] which means 'without a mind'. It is a broad conception of various gradual illnesses that consequentially interferes with one's cognition leading to behavioural issues such as inability toward self-care and daily maintenance [51]. The World Health Organisation in 2012, had announced dementia as a leading public health issue due to accelerating prevalence with population ageing. The global incidence of dementia had been estimated at 46 million people, a number that is expected to triple to 131.5 million people by the year 2050, on top of prevailing medical expenses reportedly calculated at US\$818 billion [132]. The financial strain is felt worldwide as dementia is expected to become a trillion-dollar health condition. In 2018, dementia had met the predictions made a few years prior, costing the global economy over US\$ 1 trillion with dementia being the 7th leading cause of death worldwide.

Behavioural and Psychological symptoms of dementia (BPSD), particularly impaired cognition and perception, recurrent mood shifts, and agitation [52], have posed dreadful impacts to either the person with dementia (PWD) or caregiver. The standard of living and ability to function could be diminished for the PWD [53], while BPSD could also render the role of caregiving

unbearable and depressing [53-56]. Studies also reveal in clinical settings, the motives of PWD are commonly misunderstood as destructive behaviours [57], when in fact, unfamiliar environments and people pose a threat to PWDs during their hospital visit. Therefore, to improve patient centric care, specialty training and resources must be made available for health care workers to render appropriate psychosocial strategies in crisis interventions [58-60].

End-of-Life Care

Death is the final stage in our life course process. In the ED, healthcare workers have become accustomed to death daily, despite their valiant life-saving measures. Nevertheless, end-of-life (EOL) care is seldom regarded as a fundamental knowledge component of emergency care. By the course of their practice, with maximum active resuscitation strategies, many emergency physicians are conditioned to cope with acute and severe illnesses in the ED. Nonetheless, this might be inappropriate in the management of critically ill patients experiencing impending death with a low survival rate [61,62].

Quality EOL care recognises the intricate worldview perspectives of dying, which promotes holistic patient centric care and familial support by paying attention to their physical, psychosocial, and spiritual requests for comfort [Izume et al. 2012]. In doing so, assurance of a “good death” follows regarding respect of the dying which upheld the ethical excellence in palliative care [63]. The demand for quality EOL care has become more significant with ageing population and increasing prevalence of co-morbidities [64].

Data Sources

To inform this discussion a general keyword search was undertaken to locate appropriate literature from 2010-2020. The keywords included: ‘nursing’, ‘emergency department’, ‘gerontology’, ‘geriatric nursing’, ‘geriatric emergency’, ‘emergency nursing’, ‘geriatric emergency unit’. The search aimed to identify relevant literature to create a discussion rather than perform a comprehensive and exhaustive search. Databases searched were CINAHL, MEDLINE, Google Scholar, ProQuest, ScienceDirect, and Wiley.

Discussion

It is imperative to understand that the special care needs for our older adults are generally not the primary complaints that they would present within the ED. Although the ED is intended to handle the chronically ill or wounded, it is also a sub-optimal setting for the treatment of the frail, older individual [65,66]. Previous research suggests that older adults who present to the ED with an acute illness or an exacerbation of a chronic condition have complicated medical needs, and physicians and nurses who oversee their initial treatment frequently have little expertise in geriatric medicine [67,68]. Multiple studies have shown that this lack of expertise leads to the under-triage of the seniors, compounded with increased rates of admission, delayed or disorganised treatment, increased waiting times in the ED, unnecessary diagnostic procedures, and invasive interventions [15,69-77,133]. To mitigate these shortcomings, different types of scales can be incorporated into the triage questionnaire. With consideration of quick turnovers at triage, short and validated scales should be adapted to highlight seniors at risk of functional decline.

Safety Implications

Older patients are at an increased risk for several specific events during their ED visit as compared to their younger counterparts. Impaired hearing and vision will often limit site awareness and adaptation of the older patient to an unfamiliar environment. Similarly, baseline confusion, delirium, poor balance, and spatial awareness increases the risk of injuries to the older adult in the typically designed healthcare facility [7,78]. Safety implications included hazards associated with flooring [79- 83], lighting illumination [84], problems with inpatient bathrooms [85], use of therapeutic lighting to reduce agitation [86], and problem-solving activities supported by wayfinding [87-89].

To expand ED based services, the creation of ‘geriatric emergency departments’ and ‘senior emergency rooms’ were aimed at senior citizens within ages 65 years old and above [65]. For EDs that are looking to modify their environment, they can seek help with the use of the Geriatric Emergency Department Accreditation (GEDA) guidelines developed by the American College of Emergency Physicians, Emergency Nurses Association, American Geriatrics Society, and the Society for Academic Emergency Medicine in 2013. GEDA details the changes that can be made to improve the environment for EDs to make it more geriatric-friendly. However, what it lacks is the service delivery model, which, would differ across all EDs, given the various cultural perspectives of ageing and emergency care. Thus, EDs globally would need to widely consider their ability to not only restructuring their environment but their care processes as well.

Harm was also implied when noise interrupted sleep to diminish healing and well-being [90]. The overall configuration and conditions of overcrowding created physical space limitations that were deemed unsafe [91]. This included the use of acceptable furniture/equipment to support function. For example, reclining chairs to reduce pain [92,93] and proper chair height to promote standing from sitting safely [94]. Shock-absorbing flooring [81] and grab bar placement [95] may decrease the impact of injury in falls. The use of art may yield a sense of calm [96]. Artwork of landmarks should be used in the EDs to help in the orientation of location. Older adults will feel a heightened sense of calm when they recognise landmarks of their country. Evidence has shown us that cost-effective measures can greatly impact the senior patients’ wellbeing throughout their stay, such as allowing a caregiver by his or her bedside.

Crowding as described by the American College of Emergency Physicians [97] is a circumstance when the need for emergency services outweigh possible ED or hospital reserves for patient care. Crowding, an unsought circumstance could consequentially jeopardise standards of patient care causing detrimental effects on patients’ well-being [98-100]. Backlogs in inpatient transfer to the wards after ED triage and assessment seem to be a primary feature of ED overcrowding, contributing to patients’ overstay in the ED [101]. Additional influences, such as the advent of improved healthcare needs, an aging society, a growing proportion of patients with complexities, and the emergence of advanced diagnostic technologies, can also lead to ED overcrowding [102].

No matter the circumstance, ED overcrowding may result in unfavorable patient outcomes, such as potential delay to treatment and diagnosis and higher fatality for patients admitted to inpatient and those outpatients discharged from the ED [103,104]. Functioning in stressful circumstances may jeopardise public and staff safety. Also, hastening staff “burn-out”, fortifying potential dissension amongst staff, and jeopardising the quality of care [105-108,134].

Implementation of Comprehensive Geriatric Assessment:

A comprehensive geriatric assessment (CGA) was described as a multi-dimensional interdisciplinary diagnostic process aimed at evaluating the medical, psychosocial, and functional capacities of the frail senior to establish an organised and integrative treatment plan for continuity [109]. Also, several researchers had suggested early detection of geriatric diseases using CGA in targeted patient demographics may minimise adverse outcomes pre and post hospitalisation [110-112]. Early detection of geriatric diseases can also lead to improved function and cognition capacities following hospitalisation [112-114]. When we look back at the special care needs of our seniors, we must understand that their complaints are more than meets the eye.

There are many different scales that we can use to achieve a thorough CGA which can begin at the point of triage. Firstly, a measure of functional independence. We recommend the use of the Katz Index which was created as a generalized numerical scale in order to assess, provide care, create a diagnosis, identify functional dependence in older people and people with multimorbidities [115,116]. Along with the Barthel Index, it has been widely used for ADL evaluation of patients regardless of their medical condition. Hartigan [115] and Roedel, Wilson and Fine [117] have described the Katz Index as the most appropriate scale to assess patients’ ability to perform ADL independently. The Katz Index has been chosen over the Barthel Index as it was initially developed for use in the assessment of patients with neuromuscular and musculoskeletal disorders, however, its use spread to the assessment of functional changes for patients undergoing rehab as well.

Secondly, the measure of frailty. The Clinical Frailty Scale (CFS) is a well-validated frailty measurement that originated from Dalhousie University in Canada [118]. It is scored on a scale from 1 (very fit) to 9 (terminally ill) and is based on clinical judgement [118]. Every one of the 9 levels of scores highlights a detailed explanation of the severity of the patients frailty which is accompanied by a visual aid. A score ≥ 5 is considered to be frail [118]. Information of CFS can be extracted through data from the patient’s medical charts, and therefore can also be derived from CGAs. The CFS has been validated as an adverse outcome predictor in hospitalised older people [119,120]. Simply, the CFS summarises the total fitness or frailty of a senior once they had been assessed by a competent healthcare worker [118]. Its clinical use had been further augmented to screen for frailty as it is not a questionnaire of sorts but a judgement of an encounter with the senior to report their overall health status.

Thirdly, a fall risk assessment which is currently being performed at the point of triage for all patients’ in the ED. Most EDs incorporate various measures in which they would undertake fall risk assessments. Lastly, patients undergoing palliative care. With anticipation for numerous EOL dialogues required in the ED, communication skills training is necessary for swift and effective dialogues with distraught relatives with no previous rapport built [135].

In such difficult situations, the aforementioned assessment tools would be harder to perform. However, a proper assessment will promote patient’s comfort during his or her stay in the A&E.

A comprehensive geriatric assessment of older adults is just one way to improve their health outcomes in the EDs. With quick turnovers, CGAs can be time consuming thus its assessment can be broken up into multiple layers, such as one assessment at triage and the next with a physician. The information can be further combined to develop an appropriate plan of care for the older adult.

Implications to Nursing

As healthcare providers, we need to understand that there is more beyond the scope of a patient’s complaint. The seniors that present to our ED may have an underlying medical history of dementia, frailty, functional decline, and an EOL status which could potentially be ignored. We need to be able to understand the complexity of their issues to provide appropriate treatment through different frameworks such as person-centered care or the biopsychosocial model. As such, the special care needs of older adults reiterate the importance of the environment and the role it plays in their well-being. These are also opportunities for healthcare workers to further upgrade themselves to adapt to the demands of changing demographics of our healthcare landscape to broaden their care processes in geriatrics and move away from former roles that make them task orientated.

The initial triage carried out at the “front door” of the acute hospital setting is a specialist process, which is developed for the rapid prioritization and management of life- threatening conditions. For older people, however, this triage has failed, in the past, to identify the many complexities which precipitate their admission [122]. Assessment of cognitive impairment, functional problems, and existing home care must be essential information for older adults in the ED [14,92,122].

Typical emergency department planning focuses on rapid patient assessment, evaluation, and turnover. Patient’s privacy and comfort are forsaken for greater staff manoeuvrability and flexible capacity. As such, the traditional model of emergency medical care may no longer suit the growing needs of the silver generation who seek treatment in Singapore’s ED yearly. This demographic shift brings about more challenges to our healthcare system as older people are likely to seek treatment for their multiple acute or chronic conditions. As of now, there are no workflows or processes that differentiate older adult patients from their younger counterparts. Patients presenting to the ED are triaged based on their presenting symptoms regardless of the patient’s age or functionality and our current focus is solely based on their chief complaints.

In-house training opportunities for our healthcare workers who are keen on upgrading themselves to appropriately care for our ageing population and our ageing patients’ should be conducted. Education such as lifting and transferring would also go a long way in reducing injuries that healthcare workers could sustain while caring for an older adult. Besides the older adults at risk of falling in the bathrooms, [123], healthcare workers that aid these individuals in their daily activities are at risk of injuries as well. Due to these, problems with posturing due to lifting and transferring are common in which, back pain usually follows [124,125]. This could also be a platform for

re- designing and re-skilling older workers' job scopes who are nearing their retirement. Modified job scopes can allow these older groups of healthcare workers to provide teachings for both staff and the general public to enhance their awareness of the special care needs of our senior patients.

Conclusion

With our rapidly increasing ageing population globally, we can expect an influx of our older patients both from home and long-term care facilities to present to the emergency department with a wide range of geriatric emergencies. By being able to adapt and understand their care process, we can improve patients' health outcomes and provide a quicker transition of care. However, we also do need to consider the appropriate model of care in our service delivery towards our senior patients. Research has yet to show a validated service delivery model for senior patients seeking treatment in the emergency departments. By being able to create a geriatric screening process and tailored care models, healthcare workers will be able to understand their care process and in turn, improve patients' health outcomes and provide a quicker transition of care. Further research can be performed to find solutions in improving patients' health outcomes and their service experience in the emergency department.

Declarations

Ethics Approval and Consent to Participate

Ethics approval was not required for this systematic review.

Consent for Publication

Not applicable

Availability of Data and Materials

Data sharing is not applicable to this article as no datasets were generated or analysed during the current study.

Competing Interests

The authors declare that there is no competing interests.

Funding Statement

There was no funding required for this article.

Authors' Contributions

All authors had significant contribution to this paper and meet at least one of the following criteria; (i) conception and design, or analysis and interpretation of data; (ii) drafting the article or revising it critically for important intellectual content; and (iii) final approval of the version to be published. GC, SL and MS had conceptualized and designed this discussion. GC drafted the article, which was reviewed by SL and MS. All authors approved the final version for submission.

Acknowledgements

We would like to thank Ma'am Sanda Thangarajoo and Mr. Shashi S/O Chandra Segaram for allowing our team to explore the concept of geriatric care in Singapore General Hospital's Department of Emergency Medicine.

Their guidance and feedback have allowed the visualization of a Geriatric Emergency Unit. They have given our team the utmost focus in penning down our work, stimulating our ideas and being supportive. We are very appreciative of getting the opportunity to work alongside mentors of such high calibre.

We would also like to thank Dr. Kenneth Tan Boon Kiat for his wonderful time in listening to our presentations and being supportive of our team. He has given us the confidence to pursue our research further with generous amounts of feedback as well.

Lastly, we would like to thank our nursing clinicians in Singapore General Hospital's Department of Emergency Medicine for their guidance and support through this time. They have been strong pillars leading up to this very moment. We have learnt a lot from them, and take away only good memories, new learning experiences and insights.

References

- 1) Sander M, Oxlund B, Jespersen A, Krasnik A, Mortensen EL, Westendorp RGJ, et al. The challenges of human population ageing. *Age Ageing*. 2015, 44: 185-187.
- 2) Barnett K, Mercer S, Norbury M, Watt G, Wyke S, Guthrie B. The epidemiology of multimorbidity in a large cross-sectional dataset: implications for health care, research and medical education. *Lancet*. 2012, 380: 37-43.
- 3) Salive ME. Multimorbidity in older adults. *Epidemiol Rev*. 2013, 35: 75-83.
- 4) Aminzadeh F, Dalziel WB. Older adults in the emergency department: a systematic review of patterns of use, adverse outcomes, and effectiveness of interventions. *Annals of Emergency Medicine*. 2002, 39: 238-247.
- 5) Singal BM, Hedges JR, Rousseau EW, Sanders AB, Berstein E, McNamara RM, et al. Geriatric patient emergency visits. Part I: comparison of visits by geriatric and younger patients. *Ann Emerg Med*. 1992, 21: 802-807.
- 6) Salvi F, Morichi V, Grilli A, Giorgi R, De Tommaso G, Dessi-Fulgheri P. The elderly in the emergency department: a critical review of problems and solutions. *Intern Emerg Med*. 2007, 2: 292-301.
- 7) Gray LC, Peel NM, Costa EP, Burkett E, Dey AB, Jonsson PV, et al. Profiles of older patients in the emergency department: findings from the interRAI Multinational Emergency Department Study. *Ann Emerg Med*. 2013, 62: 467-474.
- 8) Han JH, Wilber ST. Altered mental status in older emergency department patients. *Clin Geriatr Med*. 2013, 29: 101-136.
- 9) Griffiths D, Morphet J, Innes K, Crawford K, Williams A. Communication between residential aged care facilities and the emergency department. A review of the literature. *Int J Nurs Stud*. 2014, 51: 1517-1523.
- 10) Kessler C, Williams MC, Moustoukas JN, Pappas C. Transitions of care for the geriatric patient in the emergency department. *Clin Geriatr Med*. 2013, 29: 49-69.
- 11) Carpenter CR, DesPain B, Keeling TN, Shah MS, Rothenberger M. The six-item screener and AD8 for the detection of cognitive impairment in geriatric emergency department patients. *Ann Emerg Med*. 2011, 57: 653-661.

- 12) Carpenter CR, Griffey RT, Stark S, Coopersmith CM, Gage BF. Physician and nurse acceptance of technicians to screen for geriatric syndromes in the emergency department. *Western J Emerg Med*. 2011, 12: 489-495.
- 13) Carpenter CR, Heard K, Wilber S, Ginde AA, Stiffler K, Gerson LW, et al. Research priorities for high-quality Geriatric emergency care: medication management, screening, and prevention and functional assessment. *Acad Emerg Med*. 2011, 18: 644-654.
- 14) Carpenter CR, Shah MN, Hustey FM, Heard K, Gerson LW, Miller DK. High yield research opportunities in geriatric emergency medicine: prehospital care, delirium, adverse drug events, and falls. *J Gerontol A Biol Sci Med Sci*. 2011, 66: 775-783.
- 15) Schnitker L, Martin-Khan M, Beattie E, Gray L. Negative health outcomes and adverse events in older people attending emergency departments: a systematic review. *Australasian Emerg Nursing J*. 2011, 14: 141-162.
- 16) Gill TM, Allore H, Guo Z. Restricted activity and functional decline among community-living older persons. *Archives Intern Med*. 2003, 163: 1317-1322.
- 17) Katz S, Down TD, Cash HR, Grotz RC. Progress in the development of the index of ADL. *The Gerontologist*. 1970. 10: 20-30.
- 18) Lawton MP, Brody EM. Assessment of older people: self-maintaining and instrumental activities of daily living. *Gerontologist*. 1969, 9: 179-186.
- 19) Asplund K, Gustafson Y, Jacobsson C, Bucht G, Wahlin A, Peterson J, et al. Geriatric-based versus general wards for older acute medical patients: a randomized comparison of outcomes and use of resources. *J Am Geriatr Soc*. 2000, 48: 1381-1388.
- 20) Covinsky KE, Palmer RM, Fortinsky RH, Counsell SR, Stewart AL, Kresevic D, et al. Loss of independence in activities of daily living in older adults hospitalized with medical illnesses: increased vulnerability with age. *J Am Geriatr Soc*. 2003, 51: 451-458.
- 21) Boyd CM, Landefeld CS, Counsell SR, Palmer RM, Fortinsky RH, Kresevic D, et al. Recovery of activities of daily living in older adults after hospitalization for acute medical illness. *J Am Geriatr Soc*. 2008, 56: 2171-2179.
- 22) Creditor MC. Hazards of hospitalization of the elderly. *Ann Intern Med*. 1993, 118: 219-223.
- 23) Inouye SK, Wagner DR, Acampora D, Horwitz RI, Cooney LM, Hurst LM, et al. A predictive index for functional decline in hospitalized elderly medical patients. *J Gen Intern Med*. 1993, 8: 645-652.
- 24) Lee SJ, Lindquist K, Segal MR, Covinsky KE. Development and validation of a prognostic index for 4-year mortality in older adults. *JAMA*. 2006, 295: 801-808.
- 25) McCusker J, Kakuma R, Abrahamowicz M. Predictors of functional decline in hospitalized elderly patients: a systematic review. *J Gerontol A Biol Sci Med Sci*. 2002, 57: 569-577.
- 26) Rothschild JM, Bates DW, Leape LL. Preventable medical injuries in older patients. *Arch Intern Med*. 2000, 160: 2717-2728.
- 27) English KL, Paddon-Jones D. Protecting muscle mass and function in older adults during bed rest. *Curr Opin Clin Nutr Metab Care*. 2010, 13: 34-39.
- 28) Brown CJ, Redden DT, Flood KL, Allman RM. The underrecognized epidemic of low mobility during hospitalization of older adults. *J Am Geriatr Soc*. 2009, 57: 1660-1665.
- 29) Covinsky KE, Pierluissi E, Johnson B. Hospitalization associated disability "She was probably able to ambulate, but I'm not sure". *JAMA*. 2011, 306: 1782-1793.
- 30) Lakhan P, Jones M, Wilson A, Courtney M, Hirdes J, Gray LC. A prospective cohort study of geriatric syndromes among older medical patients admitted to acute care hospitals. *J Am Geriatr Soc*. 2011, 59: 2001-2008.
- 31) Landefeld CS. Care of hospitalized older patients: Opportunities for hospital-based physicians. *J Hosp Med*. 2006, 1: 42-47.
- 32) Zisberg A, Shadmi E, Sinoff G, Gur-Yaish N, Sruлович E, Admi H. Low mobility during hospitalization and functional decline in older adults. *J Am Geriatr Soc*. 2011, 59: 266-273.
- 33) Woods NF, LaCroix AZ, Gray SL, Aragaki A, Cochrane BB, Brunner RL, et al. Frailty: emergence and consequences in women aged 65 and older in the Women's Health Initiative Observational Study. *J Am Geriatr Soc*. 2005, 53: 1321-1330.
- 34) Sternberg SA, Wershof Schwartz A, Karunanathan S, Bergman H, Mark Clarfield A. The identification of frailty: a systematic literature review. *J Am Geriatr Soc*. 2011, 59: 2129-2138.
- 35) Avila-Funes JA, Carcaillon L, Helmer C, Carriere I, Ritchie K, Rouaud O, et al. Is frailty a prodromal stage of vascular dementia? Results from the Three-City Study. *J Am Geriatr Soc*. 2012, 60: 1708-1712.
- 36) Kalyani RR, Varadhan R, Weiss CO, Fried LP, Cappola AR. Frailty status and altered glucose-insulin dynamics. *J Gerontol A Biol Sci Med Sci*. 2012, 67: 1300-1306.
- 37) Gobbens RJJ, Luijckx KG, Wijnen-Sponselee MT, Schols JMGA. In search of an integral conceptual definition of frailty: opinions of experts. *J Am Med Dir Assoc*. 2010, 11: 338-343.
- 38) Gobbens RJ, Luijckx KG, Wijnen-Sponselee MT, Schols JM. Towards an integral conceptual model of frailty. *J Nutr Health Aging*. 2010, 14: 175-181.
- 39) Latham LP, Ackroyd-Stolarz S. Emergency Department Utilization by Older Adults: A Descriptive Study. *Can Geriatr J*. 2014, 17: 118.
- 40) Khandelwal D, Goel A, Kumar U, Gulati V, Narang R, Dey AB. Frailty is associated with longer hospital stay and increased mortality in hospitalized older patients. *J Nutr Health Aging*. 2012, 16: 732-735.
- 41) Kahlon S, Pederson J, Majumdar SR, Belga S, Lau D, Fradette M, et al. Association between frailty and 30-day outcomes after discharge from hospital. *CMAJ*. 2015, 187: 799-804.
- 42) Castro VM, McCoy TH, Cagan A, Rosenfield HR, Murphy SN, Churchill SE, et al. Stratification of risk for hospital admissions for injury related to fall: cohort study. *BMJ*. 2014, 349: 5863.
- 43) Ayoung-Chee P, McIntyre L, Ebel BE, Mack CD, McCormick W, Maier RV. Long-term outcomes of ground-level falls in the elderly. *J Trauma Acute Care Surg*. 2014, 76: 498-503.
- 44) Bloch F, Jegou D, Dhainaut JF, Rigaud AS, Coste J, Lundy JE, et al. Do ED staffs have a role to play in the prevention of repeat falls in elderly patients? *Am J Emerg Med*. 2009, 27: 303-307.
- 45) Denking MD, Lukas A, Nikolaus T, Hauer K. Factors associated with fear of falling and associated activity restriction in community-dwelling older adults: a systematic review. *Am J Geriatr Psychiatry*. 2015, 23: 72-86.
- 46) Evans JG. Fallers, non-fallers and Poisson. *Age Ageing*. 1990, 19: 268-269.

- 47) Murphy SL, Dubin JA, Gill TM. The development of fear of falling among community-living older women: predisposing factors and subsequent fall events. *J Gerontol A Biol Sci Med Sci.* 2003, 58: 943-947.
- 48) Davies AJ, Kenny RA. Falls presenting to the accident and emergency department: types of presentation and risk factor profile. *Age Ageing.* 1996, 25: 362-366.
- 49) Khan SA, Miskelly FG, Platt JS, Bhattacharyya BK. Missed diagnoses among elderly patients discharged from an accident and emergency department. *J Accid Emerg Med.* 1996, 13: 256-257.
- 50) Gustafson L. What is dementia? *Acta Neurologica Scandinavica.* 1996, 168: 22-24.
- 51) World Health Organization. Global action plan on the public health response to dementia 2017-2025. Geneva: World Health Organization. Licence: CC BY-NC-SA 3.0 IGO. 2017.
- 52) Finkel SI, Burns A. BPSD Consensus Statement. International Psychogeriatric Association. 1999.
- 53) Klapwijk MS, Caljouw MA, Pieper MJ, van der Steen JT, Achterberg WP. Characteristics associated with quality of life in long-term care residents with dementia: A cross-sectional study. *Dement Geriatr Cogn Disord.* 2016, 42: 186-197.
- 54) Gaugler JE, Wall MM, Kane RL, Menk JS, Sarsour K, Johnston JA, et al. The effects of incident and persistent behavioral problems on change in caregiver burden and nursing home admission of persons with dementia. *Medical Care.* 2010, 48: 875-883.
- 55) Porter CN, Miller MC, Lane M, Cormman C, Sarsour K, Kahle-Wroblewski K. The influence of caregivers and behavioral and psychological symptoms on nursing home placement of persons with Alzheimer's disease: A matched case-control study. *SAGE Open Med.* 2016, 4: 2050312116661877.
- 56) Zimmerman S, Williams CS, Reed PS, Boustani M, Preisser JS, Heck E, et al. Attitudes, stress, and satisfaction of staff who care for residents with dementia. *Gerontologist.* 2005, 1: 96-105.
- 57) Jensen AM, Pedersen BD, Olsen RB, Wilson RL, Hounsgaard L. "If only they could understand me!" Acute hospital care experiences of patients with Alzheimer's disease. *Dementia.* 2020, 19: 2331-2353.
- 58) Handley M, Bunn F, Goodman C. Supporting general hospital staff to provide dementia sensitive care: A realist evaluation. *Int J Nurs Stud.* 2019, 96: 61-71.
- 59) Hung L, Son C, Hung R. The experience of hospital staff in applying the Gentle Persuasive Approaches to dementia care. *J Psychiatr Ment Health Nurs.* 2018, 26: 19-28.
- 60) Mohler R, Meyer G. Attitudes of nurses towards the use of physical restraints in geriatric care: A systematic review of qualitative and quantitative studies. *Int J Nurs Stud.* 2014, 51: 274-288.
- 61) Grudzen CR, Richardson LD, Morrison M, Cho E, Morrison RS. Palliative care needs of seriously ill, older adults presenting to the emergency department. *Acad Emerg Med.* 2010, 17: 1253-1257.
- 62) Lamba S. Early goal-directed palliative therapy in the emergency department: a step to move palliative care upstream. *J Palliat Med.* 2009, 12: 767.
- 63) Emanuel EJ, Emanuel LL. The promise of a good death. *Lancet.* 1998, 351: S1121-S1129.
- 64) Quest TE, Marco CA, Derse AR. Hospice and palliative medicine: new subspecialty, new opportunities. *Ann Emerg Med.* 2009, 54: 94-102.
- 65) Hwang U, Morrison RS. The geriatric emergency department. *J Am Geriatr Soc.* 2007, 55, 1873-1876.
- 66) Wilber ST, Gerson LW, Terrell KM, Carpenter CR, Shah MN, Heard K, et al. Geriatric emergency medicine and the 2006 institute of medicine reports from the committee on the future of emergency care in the U.S. Health system. *J Soc Acad Emerg Med.* 2006, 13: 1345-1351.
- 67) Briggs R, Coughlan T, Collins R, O'Neill D, Kennelly SP. Nursing home residents attending the emergency department: clinical characteristics and outcomes. *QJM.* 2013, 106: 803-808.
- 68) Kennedy M, Enander RA, Tadir SP, Wolfe RE, Shapiro NI, Marcantonio ER. Delirium risk prediction, healthcare use and mortality of elderly adults in the emergency department. *J Am Geriatr Soc.* 2014, 62: 462-469.
- 69) Crilly J, Wallis M. Continuity of care for acutely unwell older adults from nursing homes. *Scand J Caring Sci.* 2006, 20: 122-134.
- 70) Crilly J, Chaboyer W, Wallis M, Thalib L, Green D. Predictive outcomes for older people who present to the emergency department. *Australasian Emerg Nurs J.* 2008, 11: 178-183.
- 71) Mudge AM, Denaro CP, O'Rourke P. Improving hospital outcomes in patients admitted from residential aged care: results from a controlled trial. *Age and Ageing.* 2012, 41: 670-673.
- 72) Hastings SN, Heflin MT. A systematic review of interventions to improve outcomes for elders discharged from the emergency department. *Acad Emerg Med.* 2005, 12: 978-986.
- 73) McCusker J, Verdon J, Tousignant P, de Courval L, Dendukuri N, Belzile E. Rapid emergency department intervention for older people reduces risk of functional decline: results of a multicenter randomized trial. *J Am Geriatr Soc.* 2001, 49: 1272-1281.
- 74) Dwyer R, Gabbe B, Stoelwinder JU, Lowthian J. A systematic review of outcomes following emergency transfer to hospital for residents of aged care facilities. *Age Ageing.* 2014, 43: 759-766.
- 75) Ackroyd-Stolarz S, Read GJ, Mackinnon NJ, Kovacs G. The association between a prolonged stay in the emergency department and adverse events in older patients admitted to hospital: a retrospective cohort study. *BMJ Qual Saf.* 2011, 20: 564-569.
- 76) Coleman EA. Falling through the cracks: challenges and opportunities for improving transitional care for persons with continuous complex care needs. *J Am Geriatr Soc.* 2003, 51: 549-555.
- 77) Coleman EA, Boulton C. American Geriatrics Society Health Care Systems C. Improving the quality of transitional care for persons with complex care needs. *J Am Geriatr Soc.* 2003, 51: 556-557.
- 78) Bradshaw LE, Goldberg SE, Lewis SA, Whittamore K, Gladman JRF, Jones RG, et al. Six-month outcomes following an emergency hospital admission for older adults with co-morbid mental health problems indicate complexity of care needs. *Age Ageing.* 2013, 42: 582-588.
- 79) Doig AK, Morse JM. The hazards of using floor mats as a fall protection device at the bedside. *J Patient Saf.* 2010, 6: 68-75.
- 80) Hewawasam L. Floor patterns limit wandering of people with Alzheimer's. *Nursing Times.* 1996, 92: 41-44.
- 81) Latimer N, Dixon S, Drahotka AK, Severs M. Costutility analysis of a shock-absorbing floor intervention to prevent injuries from falls in hospital wards for older people. *Age and Ageing.* 2013, 42: 641-645.
- 82) Simpson AH, Lamb S, Roberts PJ, Gardner TN, Evans JG. Does the type of flooring affect the risk of hip fracture? *Age and Ageing.* 2004, 33: 242-246.
- 83) Warren CJ, Hanger HC. Fall and fracture rates following a change from carpet to vinyl floor coverings in a geriatric rehabilitation hospital. A longitudinal, observational study. *Clinical Rehabilitation.* 2013, 27: 258-263.

- 84) Thuesen N, Stidsen L, Kirkegaard PH, Harder H, Suenson V. Optimizing lighting design for hospital wards by defining user zones. 1st International Conference on Lighting in Engineering, Architecture and the Environment, LIGHT. 2011, 121: 199–209.
- 85) Fink N, Pak R, Battisto D. Developing a usability evaluation tool to assess the patient room bathroom. *HERD*. 2010, 3: 22-41.
- 86) Barrick AL, Sloane PD, Williams CS, Mitchell CM, Connell BR, et al. Impact of ambient bright light on agitation in dementia. *Int J Geriatr Psychiatry*. 2010, 25: 1013-1021.
- 87) Caspi E. Wayfinding difficulties among elders with dementia in an assisted living residence. *Dementia*. 2014, 13: 429-450.
- 88) Nolan BA, Mathews RM, Harrison M. Using external memory aids to increase room finding by older adults with dementia. *Am J Alzheimers Dis Other Demen*. 2001, 16: 251-254.
- 89) Yates Bolton N, Yates K, Williamson T, Newton R. Improving hospital environments for people with dementia. Listening event report. Salford, UK: University of Salford. 2012.
- 90) Aaron JN, Carlisle CC, Carskadon MA, Meyer TJ, Hill NS, Millman RP. Environmental noise as a cause of sleep disruption in an intermediate respiratory care unit. *Sleep*. 1996, 19: 707-710.
- 91) Kelley ML, Parke B, Jokinen N, Stones M, Renaud D. Senior-friendly emergency department care: An environmental assessment. *J Health Serv Res Policy*. 2011, 16: 6-12.
- 92) Wilber ST, Burger B, Gerson LW, Blanda M. Reclining chairs reduce pain from gurneys in older emergency department patients: A randomized controlled trial. *Acad Emerg Med*. 2005, 12: 119-123.
- 93) Wilber ST, Lofgren SD, Mager TG, Blanda M, Gerson LW. An evaluation of two screening tools for cognitive impairment in older adults in an emergency department. *Acad Emerg Med*. 2005, 12: 612-616.
- 94) Weiner DK, Long R, Hughes MA, Chandler J, Studenski S. When older adults face the chair-rise challenge. A study of chair height availability and height-modified chair-rise performance in the elderly. *J Am Geriatr Soc*. 1993, 41: 6-10.
- 95) Sanford JA, Megrew MB. An evaluation of grab bars to meet the needs of elderly people. *Assist Technol*. 1995, 7: 36-47.
- 96) Wallace J, Thieme A, Wood G, Schofield G, Olivier P. Enabling self, intimacy and a sense of home in dementia: An enquiry into design in a hospital setting. Paper presented at the 30th ACM Conference on Human Factors in Computing Systems, CHI, New York, 2012, 2629-2638.
- 97) American College of Emergency Physicians. Geriatric emergency department accreditation program. American College of Emergency Physicians. 2017.
- 98) Derlet RW, Richards JR. Overcrowding in the nation's emergency departments: complex causes and disturbing effects. *Ann Emerg Med*. 2000, 35: 63-68.
- 99) Gordon JA, Billings J, Asplin BR, Rhodes KV. Safety net research in emergency medicine: proceedings of the Academic Emergency Medicine Consensus Conference on "The Unraveling Safety Net". *Acad Emerg Med*. 2001, 8: 1024-1029.
- 100) Sun BC, Hsia RY, Weiss RE, Zingmond D, Liang L-J, Han W, et al. Effect of emergency department crowding on outcomes of admitted patients. *Ann Emerg Med*. 2013, 61: 605-611.
- 101) van der Linden N, van der Linden MC, Richards JR, Derlet RW, Grootendorst DC, van den Brand CL. Effects of emergency department crowding on the delivery of timely care in an inner-city hospital in the Netherlands. *Eur J Emerg Med*. 2016, 23: 337-343.
- 102) Patterson BW, Pulia MS, Ravi S, Hoonakker PLT, Schoofs HA, Wiegmann D, et al. Scope and Influence of Electronic Health Record-Integrated Clinical Decision Support in the Emergency Department: A Systematic Review. *Ann Emerg Med*. 2019, 74: 285-296.
- 103) Pines JM, Shofer FS, Isserman JA, Abbuhl SB, Mills AM. The effect of emergency department crowding on analgesia in patients with back pain in two hospitals. *Acad Emerg Med*. 2010, 17: 276-283.
- 104) Timm NL, Ho ML, Luria JW. Pediatric emergency department overcrowding and impact on patient flow outcomes. *Acad Emerg Med*. 2008, 15: 832-837.
- 105) Barishansky RM, O'Connor KE. The effect of emergency department crowding on ambulance availability. *Ann Emerg Med*. 2004, 44: 280-281.
- 106) Barrett TW, Schriger DL. Annals of emergency medicine journal Club. Emergency department crowding is associated with poor care for patients with severe pain. *Ann Emerg Med*. 2008, 51: 6-7.
- 107) Bernstein SL, Aronsky D, Duseja R, Epstein S, Handel D, Hwang U, et al. Society for Academic Emergency Medicine, emergency department crowding task force. The effect of emergency department crowding on clinically oriented outcomes. *Acad Emerg Med*. 2009, 16: 1-10.
- 108) Weber EJ, Mason S, Freeman JV, Coster J. Implications of England's four-hour target for quality of care and resource use in the emergency department. *Ann Emerg Med*. 2012, 60: 699-706.
- 109) Rubenstein LZ, Stuck AE, Siu AL, Wieland D. Impacts of geriatric evaluation and management programs on defined outcomes: overview of the evidence. *J Am Geriatr Soc*. 1991, 39: 8S-16S.
- 110) Landefeld CS, Palmer RM, Kresevic DM, Fortinsky RH, Kowal J. A randomized trial of care in a hospital medical unit especially designed to improve the functional outcomes of acutely ill older patients. *N Engl J Med*. 1995, 332: 1338-1344.
- 111) Naylor MD, Brooten D, Campbell R, Jacobsen BS, Mezey MD, Pauly MV, et al. Comprehensive discharge planning and home follow-up of hospitalized elders: a randomized clinical trial. *JAMA*. 1999, 281: 613-620.
- 112) Van Craen CK, Braes T, Wellens N, Denhaerynck K, Flamaing J, Moons P, et al. The effectiveness of inpatient geriatric evaluation and management units: a systematic review and meta-analysis. *J Am Geriatr Soc*. 2010, 58: 83-92.
- 113) Baztan JJ, Suarez-Garcia FM, Lopez-Arrieta J, Rodriguez-Manas L, Rodriguez-Artalejo F. Effectiveness of acute geriatric units on functional decline, living at home, and case fatality among older patients admitted to hospital for acute medical disorders: meta-analysis. *BMJ*. 2009, 338: b50.
- 114) Ellis G, Langhorne P. Comprehensive geriatric assessment for older hospital patients. *Br Med Bull*. 2004, 71: 45-59.
- 115) Hartigan I. A comparative review of the Katz ADL and the Barthel Index and the Functional Independence Measure as outcome measures in stroke rehabilitation: patterns of disability scale usage in clinical trials. *Int J Rehabil Res*. 2007, 2: 204-212.

- 116) Katz S, Ford AB, Moskowitz RW, Jackson BA, Jaffe MW. Studies of illness in the aged. The Index of ADL: a standardized measure of biological and psychosocial function. *JAMA*. 1963, 185: 914-919.
- 117) Roedl KJ, Wilson LS, Fine J. A systematic review and comparison of functional assessments of community-dwelling elderly patients. *J Am Association Nurse Pract*. 2016, 28: 160-169.
- 118) Rockwood K, Song X, MacKnight C, Bergman H, Hogan DB, McDowell I, et al. A global clinical measure of fitness and frailty in elderly people. *CMAJ*. 2005, 173: 489-495.
- 119) Basic D, Shanley C. Frailty in older inpatient population: using the clinical frailty scale to predict patient outcomes. *J and Health*. 2015, 27: 670-785.
- 120) Wallis SJ, Wall J, Biram RW, Romero-Ortuno R. Association of the clinical frailty scale with hospital outcomes. *QJM*. 2015, 108:943-949.
- 121) Ellis G, Marshall T, Ritchie C. Comprehensive geriatric assessment in the emergency department. *Clin Interv Aging*. 2014, 9: 2033-2043.
- 122) Healthcare Improvement Scotland. Improving the identification and management of frailty: a case study report of innovation on four acute sites in Edinburgh: NHS Scotland. 2014.
- 123) Buzink SN, Molenbroek JFM, Haagsman EM, Bruin R, de Grootthuisen T. Falls in the toilet environment: a study on influential factors. *Gerontechnol*. 2005, 4: 15-26.
- 124) Kothiyal K, Yuen TW. Muscle strain and perceived exertion in patient handling with and without a transferring aid. *Occupational Ergonomics*. 2004, 4: 185-197.
- 125) Garg A, Owen BD, Carlson B. An ergonomic evaluation of a nursing assistants' job in a nursing home. *Ergonomics*. 1992, 35: 979-995.
- 126) Izumi S, Nagae H, Sakurai C, Imamura E. Defining end-of-life care from perspectives of nursing ethics. *Nurs Ethics*. 2012, 19: 608-618.
- 127) Moskop JC, Sklar DP, Geiderman JM, Schears RM, Bookman KJ. Emergency department crowding, part 1--concept, causes, and moral consequences. *Ann Emerg Med*. 2009, 53: 605-611.
- 128) Moskop JC, Sklar DP, Geiderman JM, Schears RM, Bookman KJ. Emergency department crowding, part 2--barriers to reform and strategies to overcome them. *Ann Emerg Med*. 2009, 53: 612-617.
- 129) World Health Organization. Dementia: a public health priority. Geneva: World Health Organization. 2012.
- 130) Fried LP, Tangen CM, Walston J, Newman AB, Hirsch C, Gottdiener J, et al. Frailty in Older Adults: Evidence for a Phenotype. *J Gerontol A Biol Sci Med Sci*. 2001, 54: 146-157.
- 131) Ang GC, Low SL, How CH. Approach to falls among the elderly in the community. *Singapore Medical J*. 2020, 61: 116-121.
- 132) Prince M, Wimo AGM, Ali GC, Wu YT, Prina M. World Alzheimer Report 2015: the global impact of dementia: an analysis of prevalence, incidence, cost and trends. London: Alzheimer's Disease International. 2015.
- 133) Geelhoed GC, de Klerk NH. Emergency department overcrowding, mortality and the 4-hour rule in Western Australia. *Med J Aus*. 2012, 196: 122-126.
- 134) Moksop JC, Sklar DP, Geiderman JM, Schears RM, Bookman KJ. Emergency department, crowding, part 1 – concept, causes, and moral consequences. *Ann Emerg Med*. 2009, 53: 605-611.
- 135) Smith AK, Fisher J, Schonberg MA, Pallin DJ, Block SD, Forrow L, et al. Am I doing the right thing? Provider perspectives on improving palliative care in the emergency department. *Ann Emerg Med*. 2009, 54: 86-93.

Submit your manuscript at

<http://enlivenarchive.org/submit-manuscript.php>

New initiative of Enliven Archive

Apart from providing HTML, PDF versions; we also provide **video version** and deposit the videos in about 15 freely accessible social network sites that promote videos which in turn will aid in rapid circulation of articles published with us.