

# Correlation between P Wave Dispersion, QRS Duration & QT Dispersion in Hospital Events in Cases of Acute Coronary Syndrome

Mahmoud Fekry Hassan Hassebo, M.B. B.Ch

Visiting Resident, Faculty of Medicine, Department of Critical Care Medicine, Alexandria University

**\*Corresponding author:** Mahmoud Fekry Hassan Hassebo, M.B. B.Ch, Visiting Resident, Faculty of Medicine, Department of Critical Care Medicine, Alexandria University, Hospitals University of Alexandria, Egypt, E-mail: mahmoudhassebo@yahoo.com

**Received Date:** 29<sup>th</sup> October 2015

**Accepted Date:** 11<sup>th</sup> February 2016

**Published Date:** 15<sup>th</sup> February 2016

**Citation:** Hassebo MFH (2016) Correlation between P Wave Dispersion, QRS Duration & QT Dispersion in Hospital Events in Cases of Acute Coronary Syndrome. Enliven: Clin Cardiol Res 3(S1): 001.

**Copyright:** © 2016 Dr. Mahmoud Fekry Hassan Hassebo. This is an Open Access article published and distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

## Summary and Conclusion

The acute coronary syndromes encompass a spectrum of unstable angina to transmural myocardial infarction. The definition of acute coronary syndrome depends on the specific characteristics of each element of the triad of clinical presentation, electrocardiographic changes and biochemical cardiac markers.

This prospective study was conducted on 60 patients with acute coronary syndrome admitted to critical care units of Alexandria University Hospitals from first of January 2012 to end of August 2012. An informed consent was taken from relatives of every patient included in the study. This study was approved by ethical committee of Alexandria Faculty of Medicine.

During the observation up to 5 days, 1 patient (1.6%) developed AF. There was no significant correlation between P wave dispersion (PD) in the three studied groups & No significant relation between it & the whole complications developed in failed thrombolysis group ( $p > 0.05$ ).

PD alone couldn't predict AF in this study, as the mean was 20.0 milliseconds, which was too small to predict AF alone.

There was no significance difference between QRS duration in the three groups (unstable angina, successful thrombolysis and failed thrombolysis respectively with mean value ( $75.7 \pm 14.02$ ,  $75.2 \pm 10.88$ ,  $73.36 \pm 8.81$ ) ( $p > 0.05$ ) & the relationship between the QRS duration and the development of complications was non-significant in this study.

This study showed that there was a significant relation among the three groups & highly significant correlation between QTD duration and prediction of complications.

It was clear that ICU length of stay was longer in patients with failed thrombolysis in comparison with the other two groups ( $p < 0.05$ ).

## Recommendations

1. Importance of measurements of QT Dispersion to patients with acute coronary syndrome especially at admission.
2. Failed thrombolysis patients should be under close observation and monitoring till another option is available like coronary intervention.

Submit your manuscript at

<http://enlivenarchive.org/submit-manuscript.php>

New initiative of Enliven Archive

Apart from providing HTML, PDF versions; we also provide **video version** and deposit the videos in about 15 freely accessible social network sites that promote videos which in turn will aid in rapid circulation of articles published with us.