

Angiotensin (1-7) Antagonist Diminished the Anti-Tumor Effect of Olmesartan in Tumor Cell Lines Grown *In-vitro* and *In-vivo*

Mohammad M. Abd-Alhaseeb^{1*}, Sawsan A. Zaitone², Soad H. Abou-El-Ela³, and Yasser M. Moustafa²

¹Department of Pharmacology and Toxicology, Faculty of Pharmacy and Pharmaceutical Industries, Sinai University, Arish, Egypt

²Department of Pharmacology and Toxicology, Faculty of Pharmacy, Suez Canal University, Ismailia, Egypt

³Department of Biochemistry, Faculty of Pharmacy and Pharmaceutical Industries, Sinai University, Arish, Egypt

*Corresponding author: Mohammad M. Abd-Alhaseeb, Department of Pharmacology and Toxicology, Faculty of Pharmacy and Pharmaceutical Industries, Sinai University, Arish, Egypt, Tel: 002-010-07699126; Fax: 002-068-3336847; E-mail: m.abdelhaseeb@su.edu.eg

Citation: Abd-Alhaseeb MM, Zaitone SA, Abou-El-Ela SH, Moustafa YM (2014) Angiotensin (1-7) Antagonist Diminished the Anti-Tumor Effect of Olmesartan in Tumor Cell Lines Grown *In-vitro* and *In-vivo*. Enliven: Challenges Cancer Detect Ther 1(1): 003.

Copyright: © 2014 Dr. Mohammad M. Abd-Alhaseeb. This is an Open Access article published and distributed under the terms of the Creative Commons Attribution License, that permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Received Date: 06th August 2014

Accepted Date: 08th September 2014

Published Date: 12th September 2014

Abstract

Olmesartan is a selective angiotensin II type 1 receptor (AT1R) antagonist. It achieves blood pressure reduction in a dose-dependent manner through arterial vasodilation and reduced sodium retention. Secondly, olmesartan exhibits anti-angiogenic activity through inhibition of Insulin growth factor, vascular endothelial growth factor and their receptors and this effect was mediated through the Ang (1-7). The current study was to investigate the anti-tumor effect of olmesartan; first, the cytotoxic activity of olmesartan and/or Ang (1-7) antagonist on MCF-7 cell line using 3-[4, 5-dimethylthiazole-2-yl]-2, 5-diphenyltetrazolium bromide (MTT) assay was explored. Then EAC solid tumor grown *in vivo* was employed to determine the impact of concurrent administration of an Ang (1-7) agonist or antagonist on the anti-tumor effect of olmesartan. In addition, the impact of concurrent administration of olmesartan on the cytotoxic activity of sorafenib in MCF-7 cell line or its anti-tumor effect in EAC solid tumor grown *in vivo* was investigated. It was observed that the cell viability was reduced by approximately 40% after sorafenib (250 µg/ml) treatment. On the other hand, olmesartan did not show any cytotoxic effect except when higher concentrations were used. IC₅₀ value for sorafenib in MCF-7 was 250.9 µg/ml, while the IC₅₀ value for olmesartan was 674.8 µg/ml. Ang (1-7) antagonist increased the IC₅₀ value of olmesartan from 674.8 µg/ml to 722 µg/ml. The high cytotoxic concentration of olmesartan in combination with sorafenib failed to enhance the cytotoxicity more than the sorafenib itself. Sorafenib (30 mg/kg/day), olmesartan (3, 10 or 30 mg/kg/day) or their combination significantly (P<0.05) reduced tumor volume and the relative tumor volume compared to EAC-Control group. Similarly, concurrent administration of the Ang (1-7) agonist with olmesartan (30 mg/kg) significantly (P<0.05) reduced tumor volume and the relative tumor volume compared to EAC-control group or olmesartan (30 mg/kg) group. Moreover, the administration of Ang (1-7) antagonist with olmesartan reduced the anti-tumor effect of olmesartan. In conclusion, olmesartan (30 mg/kg) possesses anti-tumor activity. This anti-tumor activity did not depend on the direct cytotoxic activity but might be attributed to antiangiogenic activity as proven in a previous work from our lab. The anti-tumor effect of olmesartan was, at least in part, mediated through the Ang (1-7) receptor. In addition, the present results showed that olmesartan (30 mg/kg) potentiated the anti-tumor effect of sorafenib.

Keywords:

Anti-tumor; Angiotensin (1-7) antagonist; Olmesartan; cancer; Cell line

Introduction

The Renin-angiotensin system (RAS) is a hormone system that is activated when the enzyme renin is released and cleaves the parent compound angiotensinogen to the decapeptide angiotensin I (Ang I). The catabolism of Ang I is a point of divergence in the system, leading to the production of the bioactive peptide hormones, angiotensin II (Ang II) and angiotensin (1-7) (Ang (1-7)). These peptide products differ in their carboxy termini which leads to counter-regulatory actions mediated by high affinity binding to distinct membrane-spanning receptors [1].

Ang (1-7) exerts its actions through a G protein-coupled receptor encoded by the mas gene [2]. Ang (1-7) appears to have an inhibitory influence on many of the events induced by Ang II [3]. Ang (1-7) has a depressor, vasodilator, apoptotic and anti-proliferative actions. Ang (1-7) was suggested to inhibit angiogenesis [4], although further investigations are needed to confirm these effects in a wider range of pathological/physiological conditions.

Ang (1-7) may be generated directly from Ang II by the enzymatic activity of angiotensin converting enzyme two (ACE₂) or from Ang I, via angiotensin (1-9), a pathway that utilizes both ACE₂ and angiotensin converting enzyme (ACE) [5]. ACE₂ was found in many tissues with high concentrations in the heart, kidney and gastrointestinal tract [6]. In addition, ACE₂ expression was reported in animal models of liver injury and in human cirrhosis and was associated with increasing plasma and tissue levels of Ang (1-7) [7].

Olmesartan is a selective angiotensin II type 1 receptor (AT1R) antagonist. It achieves blood pressure reduction in a dose-dependent manner through arterial vasodilation and reduced sodium retention [8]. In addition, olmesartan exhibits anti-angiogenic activity through inhibition of Insulin growth factor, vascular endothelial growth factor and their receptors and this effect was mediated through the Ang (1-7) [9]. Sorafenib is a multi-kinase inhibitor taken orally and approved in the treatment of metastatic renal cell carcinoma [10]. It has been reported that olmesartan potentiated the anti-angiogenic effect of sorafenib in Ehrlich's ascites carcinoma (EAC) solid tumor grown *in vivo* in mice [9]. So, the objective of the current study was to investigate the anti-tumor effect of olmesartan; beginning with exploring the cytotoxic activity of olmesartan and/or Ang (1-7) antagonist on MCF-7 cell line using 3-[4, 5-dimethylthiazole-2-yl]-2, 5-diphenyltetrazolium bromide (MTT) assay. Then, EAC solid tumor grown *in vivo* was employed to determine the impact of concurrent administration of an Ang (1-7) agonist or antagonist on the anti-tumor effect of olmesartan. Finally, investigate the impact of concurrent administration of olmesartan on the cytotoxic activity of sorafenib in MCF-7 cell line or its anti-tumor effect in EAC solid tumor grown *in vivo*.

Methods and Materials

Cell Culture and Drug Treatment

The MCF-7 human breast adenocarcinoma cell line was purchased from American Type Culture Collection (ATCC, Manassas, VA, USA). It was maintained in Dulbecco's Modified Eagle's Medium containing 10% fetal bovine serum, 100 units/mL penicillin, and 100 mg/mL streptomycin. Cells were incubated in a humidified, 5% CO₂ atmosphere at 37°C.

MTT Assay for Cell Viability

MTT assay is based on the ability of active mitochondrial dehydrogenase enzyme of living cells to cleave the tetrazolium rings of the yellow MTT and form dark blue insoluble formazan crystals which is largely impermeable to cell membranes, resulting in its accumulation within healthy cells. The effect of olmesartan and/or sorafenib and Ang (1-7) antagonist on cell viability was determined using MTT assay. In MTT assay 0.5×10⁵ cells per well were plated in 96-well culture plates. After an overnight incubation, cells were treated with 20 µl of different concentrations of olmesartan and/or sorafenib for 48 h at 37°C. The cells were then treated with 40 µl of MTT (Sigma-Aldrich, MO, USA) and incubated for 4 h at 37°C. The medium was then discarded, and 180 µl of acidified isopropanol (Sigma-Aldrich, MO, USA) was added to dissolve formazan crystals.

Absorption values at 570 nm were determined with Multiskan MS microplate reader (Labsystems, Finland). The cell viability of olmesartan and/or sorafenib-treated cells was calculated as the percentage of cell viability compared to untreated cells. In addition, IC₅₀ values were calculated from the equation of the curve.

Anti-Tumor Activity of Olmesartan and/or Sorafenib in Ehrlich's Ascites Carcinoma Solid Tumor Grown in Mice

Female Swiss albino mice, each weighing 20–25 g were obtained from the modern veterinary office for laboratory animals (Cairo, Egypt). EAC cell line was purchased from Tumor Biology Department, National Cancer Institute (Cairo University, Egypt). EAC cells were injected intradermally (2.5 × 10⁶ EAC cells in 0.1 ml saline/animal) at the two sites bilaterally on the lower ventral side after shaving this area. After 7 days, mice were randomly divided into eight groups, ten animals each.

All treatments were given for 21 days and the treatment regimens were as follows: Group I: mice treated with DMSO (5 mL/kg/day, p.o.), and served as the EAC-control group. Group II: mice treated with sorafenib (30 mg/kg/day, p.o.) [11]. Group III-V: mice treated with olmesartan (3, 10 or 30 mg/kg/day, p.o.), respectively [12]. Group VI: mice treated with a combination of sorafenib (30 mg/kg/day, p.o.) and olmesartan (30 mg/kg/day, p.o.). Group VII: mice treated with olmesartan (30 mg/kg/day, p.o.) and the angiotensin (1-7) agonist (30 µg/kg/day, i.p.) [13]. Group VIII: mice were treated with olmesartan (30 mg/kg/day, p.o.) and the angiotensin (1-7) antagonist (A-779 peptide) (3.3 mg/kg/trice weekly, i.p.) [14]. In general, olmesartan and sorafenib were administered daily by gastric gavage in a volume of 5 mL/kg. Whereas, the angiotensin (1-7) agonist or the angiotensin (1-7) antagonist were administered intraperitoneally.

At the end of the experiment, the animals were sacrificed with cervical dislocation. The tumors were separated from the surrounding muscles and dermis; tumor volumes were measured with vernier calipers and calculated by the following formula: 0.5 X²Y, where X and Y are the minor and major axes, respectively [15]. In addition, the relative tumor volumes were calculated by dividing the mean tumor volumes of the treated groups by the mean tumor volume of the control group [16]. All experimental protocols were approved by The Research Ethics Committee at the Faculty of Pharmacy, Suez Canal University (License number 20146A10).

Drugs and Chemicals

Olmesartan medoxomil was obtained from Daiichi Sankyo Pharmaceutical Co. (Tokyo, Japan) and dissolved at a concentration of 100 mM in dimethyl sulphoxide (DMSO, Sigma-Aldrich, MO, USA) as a stock solution. It was then further diluted to working concentrations with cell culture medium in *in-vitro* study and with water in *in-vivo* study. Sorafenib tosylate was purchased from Bayer Health Care (Leverkusen, Germany). Ang (1-7) agonist and antagonist were purchased from Bachem AG (Bubendorf, Zurich, Switzerland). All other chemicals were purchased from Sigma-Aldrich (MO, USA).

Statistical Analysis

In-vitro results were expressed as mean \pm standard deviation (SD). Results were analysed in terms of IC_{50} values, and differences noted across the cell-line panel and within individual cell lines were tested for statistical significance using Chi-square test. On the other hand data from *in-vivo*

results were expressed as mean \pm standard error of mean (SEM) and was analyzed using one-way analysis of variance (ANOVA), followed by Bonferroni's post hoc test at $P < 0.05$. Statistical analysis was performed using SPSS software, version 22 (SPSS Software, SPSS Inc., Chicago, USA).

Results

Effect on MCF-7 Cell Line

First, we determined the cytotoxic effect of sorafenib and olmesartan on MCF-7 breast cancer cells using MTT assay. MCF-7 cells were treated with various concentrations of sorafenib and olmesartan. After sorafenib treatment, cell viability was reduced by approximately 40% (Figure 1A). On the other hand, olmesartan did not show any cytotoxic effect except when higher concentrations were used (Figure 1B).




Figure 1: Effect of sorafenib and olmesartan on cell viability in MCF-7 cells. A) Effect of sorafenib against MCF-7 cells.

B) Effect of olmesartan against MCF-7 cells. Cell viability was measured using MTT assay. Values are expressed as mean value of cell viability (% of control) \pm S.D. of four experiments and analyzed using Chi-square test. *Significantly different from control at $P < 0.05$.

IC_{50} value for sorafenib in MCF-7 was 250.9 $\mu\text{g/ml}$, while the IC_{50} value for olmesartan was 674.8 $\mu\text{g/ml}$ (Figure 2A and 2B). On the other hand, the Ang (1-7) antagonist (A-779 peptide) showed a safe effect on the same cell line up to 50 $\mu\text{g/ml}$, so that the used concentration (10 $\mu\text{g/ml}$) was completely safe with viability percent $> 85\%$.

In the low concentration range up to 100 $\mu\text{g/ml}$ of sorafenib or olmesartan, the 10 $\mu\text{g/ml}$ of peptide decreased the IC_{50} of sorafenib from 250.9 $\mu\text{g/ml}$ to 125.9 $\mu\text{g/ml}$. On the other hand, the peptide (10 $\mu\text{g/ml}$) treatment in combination with olmesartan increased the IC_{50} value from 674.8 $\mu\text{g/ml}$ to 722 $\mu\text{g/ml}$ (Figure 3A and 3B).




Figure 2: The half maximal inhibitory concentrations of sorafenib and olmesartan in MCF-7 cells. A) IC_{50} value for sorafenib in MCF-7 cells.

B) IC_{50} value for olmesartan in MCF-7 cells. Cell viability was measured using MTT assay. IC_{50} : The half maximal inhibitory concentration.




Figure 3: Effect of angiotensin (1-7) antagonist (A-779 peptide) on IC₅₀ value of sorafenib and olmesartan using MCF-7 cells. A) Effect of sorafenib with the A-779 peptide against MCF-7 cells. B) Effect of olmesartan with A-779 peptide against MCF-7 cells. Cell viability was measured using MTT assay. IC₅₀: The half maximal inhibitory concentration.

The combination of olmesartan and sorafenib, with different concentrations of both, showed different effects (Figure 4). Olmesartan itself at highest used concentration (500 µg/ml) - without sorafenib - enhanced cytotoxicity from about 90% of cell viability - at olmesartan concentration 62.5 µg/ml - into only 60% of cell viability.

However such high cytotoxic concentration of olmesartan in combination with sorafenib failed to enhance the cytotoxicity more than the sorafenib itself (Figure 4).




Figure 4: Cytotoxic effect of fixed concentrations of olmesartan and/or sorafenib against MCF-7 cell line. Cell viability was measured using MTT assay.

Effect on Tumor Volume

Administration of sorafenib (30 mg/kg/day), olmesartan (3, 10 or 30 mg/kg/day) or their combination significantly ($P < 0.05$) reduced tumor volume compared to EAC-Control group (Figure 5A). Similarly, concurrent administration of the Ang (1-7) agonist with olmesartan (30 mg/kg) significantly ($P < 0.05$) reduced tumor volume compared to EAC-control group or olmesartan (30 mg/kg) group. Moreover, the administration of Ang (1-7) antagonist with olmesartan reduced the antitumor effect of olmesartan (Figure 5B).

Effect on the Relative Tumor Volume

The administration of sorafenib (30 mg/kg/day), olmesartan (3, 10 or 30 mg/kg/day) or their combination showed a significant ($P < 0.05$) decrease in the % relative tumor volume when compared to EAC-control group (Figure 6A). Similarly, concurrent administration of the Ang (1-7) agonist with olmesartan (30 mg/kg) significantly ($P < 0.05$) reduced the % relative tumor volume compared to EAC-Control group or olmesartan (30 mg/kg) group. Further, the administration of Ang (1-7) antagonist with olmesartan reduced the antitumor effect of olmesartan (Figure 6B).




Figure 5: Effect of sorafenib and olmesartan on tumor volume of EAC solid tumor grown in mice. A) Effect of sorafenib (30 mg/kg), olmesartan (3, 10 or 30 mg/kg) and their combination on the mean tumor volume of EAC solid tumor growing in mice. B) Effect of concurrent administration of an Ang (1-7) agonist (30 µg/kg/day, i.p.) or an Ang (1-7) antagonist (3.3 mg/kg/trice/week, i.p.) and olmesartan on the mean tumor volume of EAC solid tumor grown in mice. EAC: Ehrlich's ascites carcinoma. Values are expressed as the mean ± S.E.M. and data were analyzed using one-way ANOVA followed by Bonferroni's post-hoc test at $P < 0.05$. *Significantly different from the EAC-control. ^ΔSignificantly different from sorafenib monotherapy. [§]Significantly different from olmesartan (3 mg/kg) group. [€]Significantly different from olmesartan (10 mg/kg) group. [§]Significantly different from olmesartan (30 mg/kg) group. [◊]Significantly different from the combination of olmesartan and Ang (1-7) agonist.




Figure 6: Effect of sorafenib and olmesartan on the relative tumor volume of EAC solid tumor grown in mice. A) Effect of sorafenib (30 mg/kg), olmesartan (3, 10 or 30 mg/kg) and their combination on the relative tumor volume of EAC solid tumor grown in mice. B) Effect of concurrent administration of an Ang (1-7) agonist (30 µg/kg/day, i.p.) or an Ang (1-7) antagonist (3.3 mg/kg/trice/week, i.p.) and olmesartan on the relative tumor volume of EAC solid tumor grown in mice. EAC: Ehrlich's ascites carcinoma. Values are expressed as the mean ± S.E.M. and data were analyzed using one-way ANOVA followed by Bonferroni's post-hoc test at $P < 0.05$. *Significantly different from the EAC-control. ^ΔSignificantly different from sorafenib monotherapy. [§]Significantly different from olmesartan (3 mg/kg) group. [€]Significantly different from olmesartan (10 mg/kg) group. [§]Significantly different from olmesartan (30 mg/kg) group. [◊]Significantly different from the combination of olmesartan and Ang (1-7) agonist.

Discussion

There are increasing evidences for *in-vivo* and *in-vitro* models of angiogenesis indicating a regulatory role of Ang II and its receptors in new vessel formation [17]. Ang II has been reported to promote tumor growth and angiogenesis [18]. Therefore, angiotensin receptor blockers have been considered a noteworthy anticancer and anti-angiogenesis therapeutic option [18].

Angiotensin II type 1 receptor is often up-regulated during the progression from normal to malignant phenotypes, indicating at the very least a correlation between the RAS and tumour progression [3]. Therefore, AT₁R blockers have been considered as an anti-angiogenic therapeutic option [19].

Ang (1-7) appears to have an inhibitory influence on many of the events induced by Ang II [3]. Ang (1-7) has a depressor, vasodilator, apoptotic and anti-proliferative actions. Ang (1-7) is also suggested to inhibit angiogenesis [4].

In the current study, olmesartan showed a cytotoxic activity and reduced the cell viability of MCF-7 cells with IC₅₀ value of 675 µg/ml; however this is considered a high cytotoxic concentration. Therefore, we suggested that the anti-tumor effect of olmesartan is not a result of direct toxicity. Consistently, it has been reported that the antitumor effect of ARBs is not a result of direct toxicity but of an anti-angiogenic effect [20,21]. In addition, it has been reported that in the MCF-7 cell line, Ang II increased the basal protein kinase activity and so increased growth of MCF-7 cells. Consequently, ARBs decreased growth of MCF-7 cells through inhibition of protein kinase activity not due to cytotoxic activity [22]. Another study came in parallel with the present findings as candesartan, type of ARBs, did not induce direct cytotoxicity in *in-vitro* human bladder cancer cells [20].

In addition, the current results demonstrated that the Ang (1-7) antagonist increased the IC₅₀ value of olmesartan indicated the antagonist effect exerted by the Ang (1-7) antagonist on olmesartan. In agreement with the previous results, it has been reported that the specific Ang (1-7) receptor antagonist (A-779 peptide) prevented the effects of ARBs and Ang (1-7) itself [4].

On the other hand, sorafenib showed a higher cytotoxic activity against MCF-7 cell lines with IC₅₀ value of 250 µg/ml; that indicated the higher cytotoxicity of sorafenib over olmesartan. In agreement with the previous results, it has been reported that sorafenib showed a broad cytotoxic activity against various tumor cell lines *in-vitro* and in xenograft models [23].

Additionally, the current study showed that olmesartan at the highest used concentration (500 µg/ml) enhanced cytotoxicity from about 90% of cell viability into only 60% of cell viability. Therefore, olmesartan showed a little cytotoxic activity. In agreement with the previous results, it has been reported that the ARBs showed a mild cytotoxic activity on tumor cell lines [21].

This is the first time to examine the cytotoxic effect of olmesartan and/or sorafenib on MCF-7 cells. The current results showed that the highest cytotoxic concentration of olmesartan in combination with sorafenib failed to enhance the cytotoxicity more than the sorafenib itself indicating no *in-vitro* synergistic effect in the cytotoxicity between the two compounds despite of the toxicity of each one separately.

The *in-vivo* anti-tumor activity of olmesartan was evaluated in the present study by determination of tumor volume and the relative tumor volume in EAC solid tumor grown in mice. The current study showed that olmesartan reduced the tumor volume and relative tumor volume assuming that this was linked to the angiostatic effect of olmesartan which resulted in tumor growth impairment. In agreement with the previous results, it has been reported that candesartan reduced tumor volume in a xenograft model of bladder cancer [20].

Furthermore, consistently with the previous results losartan reduced cell growth of *in-vivo* models of cancer [24]. Also, telmisartan, caused marked inhibition of prostate cancer cells in concentration-dependent and time-dependent manner [18].

The current results showed that the combination of olmesartan (30 mg/kg) with the Ang (1-7) agonist reduced the tumor volume and the relative tumor volume. On the other hand, the Ang (1-7) antagonist (A-779 peptide) antagonized the anti-tumor effect of olmesartan. Therefore, we suggested that the anti-tumor effect of olmesartan is mediated through the Ang (1-7) receptors. In agreement with the previous results it has been reported that the Ang (1-7) antagonist antagonized the anti-tumor effect of Ang (1-7) agonist and ARBs in human lung cancer cell model [25].

The current study also showed that sorafenib reduced tumor volume and the relative tumor volume. In agreement with the previous results, it has been reported that sorafenib reduced tumor size and tumor weight in hepatocellular carcinoma [26]. In addition, it has been reported that sorafenib reduced tumor weight and tumor volume in neuroblastoma model of cancer [27]. Another study came in parallel with the results in the current study, it showed that sorafenib reduced tumor weight in human liver cancer model [28].

Moreover, the current study showed that olmesartan (30 mg/kg) potentiated the anti-tumor effect of sorafenib. The combined therapy reduced tumor volume and the relative tumor volume and this effect was attributed to the anti-angiogenic effect of the combined therapy. Similarly, it has been reported in previous study that the combined therapy of olmesartan and sorafenib produced an anti-angiogenic activity that was confirmed by reducing tumor weight of EAC solid tumor grown on mice [9].

Conclusion

In conclusion, the present results showed that olmesartan (30 mg/kg) possesses anti-tumor activity. This anti-tumor activity did not depend on the direct cytotoxic activity but might be attributed to antiangiogenic activity as proven in a previous work from our lab. The anti-tumor effect of olmesartan was, at least in part, mediated through the Ang (1-7) receptor. In addition, the present results showed that olmesartan (30 mg/kg) potentiated the anti-tumor effect of sorafenib. Therefore, the present study highlights the beneficial role of olmesartan as an adjuvant medication to sorafenib in the treatment of cancer.

References

1. Ferrario CM, Jessup J, Gallagher PE, Averill DB, Brosnihan KB, et al. (2005) Effects of renin-angiotensin system blockade on renal angiotensin-(1-7) forming enzymes and receptors. *Kidney Int* 68: 2189-2196.
2. Santos RA, Simoes e Silva AC, Maric C, Silva DM, Machado RP, et al. (2003) Angiotensin-(1-7) is an endogenous ligand for the G protein-coupled receptor Mas. *Proc Natl Acad Sci U S A* 100: 8258-8263.
3. Ager EI, Neo J, Christophi C (2008) The renin-angiotensin system and malignancy. *Carcinogenesis* 29: 1675-1684.
4. Machado RD, Santos RA, Andrade SP (2001) Mechanisms of angiotensin-(1-7)-induced inhibition of angiogenesis. *Am J Physiol Regul Integr Comp Physiol* 280: R994-R1000.
5. Herath CB, Warner FJ, Lubel JS, Dean RG, Jia Z, et al. (2007) Upregulation of hepatic angiotensin-converting enzyme 2 (ACE2) and angiotensin-(1-7) levels in experimental biliary fibrosis. *J Hepatol* 47: 387-395.
6. Donoghue M, Hsieh F, Baronas E, Godbout K, Gosselin M, et al. (2000) A novel angiotensin-converting enzyme-related carboxypeptidase (ACE2) converts angiotensin I to angiotensin 1-9. *Circ Res* 87: E1-9.
7. Paizis G, Tikellis C, Cooper ME, Schembri JM, Lew RA, et al. (2005) Chronic liver injury in rats and humans upregulates the novel enzyme angiotensin converting enzyme 2. *Gut* 54: 1790-1796.
8. Mire DE, Silfani TN, Pugsley MK (2005) A review of the structural and functional features of olmesartan medoxomil, an angiotensin receptor blocker. *J Cardiovasc Pharmacol* 46: 585-593.
9. Abd-Alhaseeb MM, Zaitone SA, Abou-El-Ela SH, Moustafa YM (2014) Olmesartan potentiates the anti-angiogenic effect of sorafenib in mice bearing Ehrlich's ascites carcinoma: role of angiotensin (1-7). *PLoS one* 9: e85891.
10. Ratain MJ, Eisen T, Stadler WM, Flaherty KT, Kaye SB, et al. (2006) Phase II placebo-controlled randomized discontinuation trial of sorafenib in patients with metastatic renal cell carcinoma. *J Clin Oncol* 24: 2505-2512.
11. Huynh H, Lee JW, Chow PK, Ngo VC, Lew GB, et al. (2009) Sorafenib induces growth suppression in mouse models of gastrointestinal stromal tumor. *Mol Cancer Ther* 8: 152-159.
12. Tsuda M, Iwai M, Li JM, Li HS, Min LJ, et al. (2005) Inhibitory effects of AT1 receptor blocker, olmesartan, and estrogen on atherosclerosis via anti-oxidative stress. *Hypertension* 45: 545-551.
13. Fraga-Silva RA, Costa-Fraga FP, De Sousa FB, Alenina N, Bader M, et al. (2011) An orally active formulation of angiotensin-(1-7) produces an antithrombotic effect. *Clinics* 66: 837-841.
14. Jawien J, Toton-Zuranska J, Gajda M, Niepsuj A, Gebaska A, et al. (2012) Angiotensin-(1-7) receptor Mas agonist ameliorates progress of atherosclerosis in apoE-knockout mice. *J Physiol Pharmacol* 63: 77-85.
15. Attia MA, Weiss DW (1966) Immunology of spontaneous mammary carcinomas in mice. V. Acquired tumor resistance and enhancement in strain A mice infected with mammary tumor virus. *Cancer Res* 26: 1787-1800.
16. Wild R, Dings RP, Subramanian I, Ramakrishnan S (2004) Carboplatin selectively induces the VEGF stress response in endothelial cells: Potentiation of antitumor activity by combination treatment with antibody to VEGF. *Int J Cancer* 110: 343-351.
17. Carbajo-Lozoya J, Lutz S, Feng Y, Kroll J, Hammes HP, et al. (2012) Angiotensin II modulates VEGF-driven angiogenesis by opposing effects of type 1 and type 2 receptor stimulation in the microvascular endothelium. *Cell Signal* 24: 1261-1269.
18. Funao K, Matsuyama M, Kawahito Y, Sano H, Chargui J, et al. (2008) Telmisartan is a potent target for prevention and treatment in human prostate cancer. *Oncology Rep* 20: 295-300.
19. Abali H, Gullu IH, Engin H, Haznedaroglu IC, Erman M, et al. (2002) Old antihypertensives as novel antineoplastics: angiotensin-I-converting enzyme inhibitors and angiotensin II type 1 receptor antagonists. *Med Hypotheses* 59: 344-348.
20. Kosugi M, Miyajima A, Kikuchi E, Horiguchi Y, Murai M (2006) Angiotensin II type 1 receptor antagonist candesartan as an angiogenic inhibitor in a xenograft model of bladder cancer. *Clinical Can Res* 12: 2888-2893.
21. Kosaka T, Miyajima A, Takayama E, Kikuchi E, Nakashima J, et al. (2007) Angiotensin II type 1 receptor antagonist as an angiogenic inhibitor in prostate cancer. *The Prostate* 67: 41-49.
22. Loewenthal H, Jahn G (1932) Übertragung-Suersuche Mit Carcinomatöser Mause-Asciteslussigkeit Und Ihr Verhalten Gegen Physikalische Und Chemische Einwirkungen. *Zeitschrift für Krebsforschung* 37: 439-447.
23. Yu C, Friday BB, Lai JP, Yang L, Sarkaria J, et al. (2006) Cytotoxic synergy between the multikinase inhibitor sorafenib and the proteasome inhibitor bortezomib *in vitro*: induction of apoptosis through Akt and c-Jun NH2-terminal kinase pathways. *Mol Cancer Ther* 5: 2378-2387.
24. Arrieta O, Guevara P, Escobar E, Garcia-Navarrete R, Pineda B, et al. (2005) Blockage of angiotensin II type I receptor decreases the synthesis of growth factors and induces apoptosis in C6 cultured cells and C6 rat glioma. *Br J Cancer* 92: 1247-1252.
25. Soto-Pantoja DR, Menon J, Gallagher PE, Tallant EA (2009) Angiotensin-(1-7) inhibits tumor angiogenesis in human lung cancer xenografts with a reduction in vascular endothelial growth factor. *Mol Cancer Ther* 8: 1676-1683.
26. Kong J, Kong F, Gao J, Zhang Q, Dong S, et al. (2014) YC-1 enhances the anti-tumor activity of sorafenib through inhibition of signal transducer and activator of transcription 3 (STAT3) in hepatocellular carcinoma. *Mol Cancer* 13: 7.
27. Kakodkar NC, Peddinti RR, Tian Y, Guerrero LJ, Chlenski A, et al. (2012) Sorafenib inhibits neuroblastoma cell proliferation and signaling, blocks angiogenesis, and impairs tumor growth. *Pediatr Blood Cancer* 59: 642-647.
28. Kusano H, Ogasawara S, Akiba J, Nakayama M, Ueda K, et al. (2013) Antiproliferative effects of sorafenib and pegylated IFNalpha2b on human liver cancer cells in vitro and in vivo. *International journal of oncology* 42: 1897-1903.

Submit your manuscript at
<http://enlivenarchive.org/submit-manuscript.php>

New initiative of Enliven Archive

Apart from providing HTML, PDF versions; we also provide **video version** and deposit the videos in about 15 freely accessible social network sites that promote videos which in turn will aid in rapid circulation of articles published with us.